



Kjeldsen, K. K., Korsgaard, N. J., Bjørk, A. A., Khan, S. A., Box, J. E., Funder, S., Larsen, N. K., Bamber, J. L., Colgan, W., Van Den Broeke, M., Siggaard-Andersen, M. L., Nuth, C., Schomacker, A., Andresen, C. S., Willerslev, E., & Kjær, K. H. (2015). Spatial and temporal distribution of mass loss from the Greenland Ice Sheet since AD 1900. *Nature*, 528(7582), 396-400.
<https://doi.org/10.1038/nature16183>

Peer reviewed version

License (if available):
Unspecified

Link to published version (if available):
[10.1038/nature16183](https://doi.org/10.1038/nature16183)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Nature at <http://www.nature.com/nature/journal/v528/n7582/full/nature16183.html>.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**Greenland freshwater pathways in the sub-Arctic Seas from model experiments with
passive tracers**

Corresponding author: Dmitry S. Dukhovskoy, Center for Ocean-Atmospheric Prediction
Studies, The Florida State University, 2000 Levy Avenue, Building A, Suite 257,
Tallahassee, FL 32310, USA, (ddukhovskoy@fsu.edu)

Paul G. Myers, Department of Earth and Atmospheric Sciences, University of Alberta, 1-26
Earth Sciences Building, University of Alberta, Edmonton, Alberta, T6G 2E3,
Canada, (pmyers@ualberta.ca)

Gennady Platov, (a) Institute of Computational Mathematics and Mathematical Geophysics,
630090, Lavrentiev prospect 6, Novosibirsk, Russia; (b) Novosibirsk State University,
Novosibirsk, Russia; (Platov.G@gmail.com)

Mary-Louise Timmermans, Department of Geology and Geophysics, Yale University, 210
Whitney Ave., New Haven CT, 06511, USA (mary-louise.timmermans@yale.edu)

Beth Curry, Applied Physics Laboratory, University of Washington, 1013NE40th St.,
Seattle, WA98105, (beth4cu@uw.edu)

Andrey Proshutinsky, Department of Physical Oceanography, Woods Hole Oceanographic
Institution, Mail Stop 29 / 218B Clark Laboratory, Woods Hole, MA 02543, USA,
(aproshutinsky@whoi.edu)

Jonathan L. Bamber, Bristol Glaciology Centre, School of Geographical Sciences,
University of Bristol, University Road, Bristol, BS8 1SS, UK,
(J.Bamber@bristol.ac.uk)

Eric Chassignet, Center for Ocean-Atmospheric Prediction Studies, The Florida State
University, 2000 Levy Avenue, Building A, Suite 252, Tallahassee, FL 32310, USA,
(echassignet@fsu.edu)

Xianmin Hu, Department of Earth and Atmospheric Sciences, 1-26 ESB, University of
Alberta, Edmonton, Alberta, T6G 2E3, Canada (xianmin@ualberta.ca)

Craig M. Lee, Applied Physics Laboratory, University of Washington, 1013 NE 40th St.,
Seattle, WA 98105, (craig@apl.washington.edu)

Raquel Somavilla, Spanish Institute of Oceanography, Avda. Principe de Asturias, 70 Bis,
33212, Gijón, Spain, (raquel.somavilla@gi.ieo.es).

Abstract

Accelerating since the early 1990s, the Greenland Ice Sheet mass loss exerts a significant impact on thermohaline processes in the sub-Arctic seas. Surplus freshwater discharge from Greenland since the 1990s, comparable in volume to the amount of freshwater present during the Great Salinity Anomaly events, could spread and accumulate in the sub-Arctic seas and influence convective processes there. However, hydrographic observations in the Labrador Sea and the Nordic Seas, where the Greenland freshening signal might be expected to propagate, do not show a persistent freshening in the upper ocean during last two decades. This raises the question of where the surplus Greenland freshwater has propagated. In order to investigate the fate, pathways, and propagation rate of Greenland freshwater in the sub-Arctic seas, several numerical experiments using a passive tracer to track the spreading of Greenland freshwater have been conducted as a part of the Forum for Arctic Ocean Modeling and Observational Synthesis effort. The models show that Greenland freshwater propagates and accumulates in the sub-Arctic seas, although there is disagreement among the models on the amount and rate of tracer propagation into the convective regions. Results highlight the differences in simulated physical mechanisms at play in different models and underscore the continued importance of intercomparison studies. It is estimated that surplus Greenland freshwater flux should have caused a salinity decrease by 0.06—0.08 in the sub-Arctic seas, which is smaller than the recently observed salinization (0.15–0.2) of Atlantic Water that has possibly obscured the freshening signal.

Key words: Greenland Ice Sheet melting, Greenland freshwater, thermohaline circulation, Nordic Seas, sub-Arctic seas, Baffin Bay, Labrador Sea

3 key points:

Results from passive tracer model experiments are presented

Greenland freshwater pathways and accumulation in the sub-Arctic seas are analyzed

Meltwater influence on salinity change in the sub-Arctic is discussed

1. Introduction

Observational and modeling studies indicate that the Greenland Ice Sheet and other Arctic land ice are melting at a rate that has dramatically increased since the early 1990s [Pritchard *et al.*, 2009; Velicogna, 2009; Gardner *et al.*, 2011; Bamber *et al.*, 2012; Box and Colgan, 2013]. The volume of surplus freshwater discharge from Greenland into the sub-Arctic seas (Greenland, Iceland, Norwegian, Labrador seas, Baffin Bay, and the Subpolar region in the North Atlantic, Figure 1) since 1990 (in 2010 $\sim 3,200 \text{ km}^3$) is approaching the magnitude of the freshwater anomaly ($10,000 \text{ km}^3$) advected into the North Atlantic during the 1970 Great Salinity Anomaly event (GSA) [Dickson *et al.*, 1998; Bamber *et al.*, 2012], suggesting that this freshwater source may have significant impact on ocean conditions in the region.

The sensitivity of the thermohaline circulation in the North Atlantic to freshwater balance has been widely discussed in the literature [Stommel, 1961; Rooth, 1982; Manabe and Stouffer, 1988; Rahmstorf, 1995; Malmberg and Jonsson, 1997; Stouffer *et al.*, 2006; Jahn and Holland, 2013]. Considerable progress has been made in understanding the role of the thermohaline circulation in abrupt climate change from the paleoclimate perspective [e.g., Manabe and Stouffer, 1993; 1995; Clark *et al.*, 2002], thus providing a link between the freshwater budget of the sub-Arctic and climate variability. The concept of freshwater influence on the thermohaline circulation has been generally accepted and used to explain observed climate variability in the Arctic and North Atlantic [Stouffer *et al.*, 2015 and references therein]. The Arctic Ocean and the Greenland Ice Sheet are two sources of freshwater that potentially may impact the thermohaline circulation in the North Atlantic and influence the Atlantic Meridional Overturning Circulation (AMOC).

Past GSA events resulting from anomalously high freshwater flux from the Arctic Ocean have been shown to have impacted thermohaline conditions and climate in the North Atlantic in the 20th century [Dickson *et al.*, 1988; Belkin, 2004; Belkin *et al.*, 2008]. Further, conceptual and idealized models have put forward mechanisms by which a freshwater flux from the Arctic Ocean may be a key factor in controlling the thermohaline circulation in the North Atlantic [e.g., Ikeda, 1990; Mysak and Venegas, 1998; Ikeda *et al.*, 2001; Goosse *et al.*, 2002; Proshutinsky *et al.*, 2002; Dukhovskoy *et al.*, 2004].

Meltwater from the Greenland Ice Sheet is the other major source of freshwater considered to influence thermohaline circulation and climate [Rahmstorf, 2002; Fichefet *et al.*, 2003; Rahmstorf, 2005; Ridley *et al.*, 2005; Rudels, 2011; Castro de la Guardia *et al.*, 2015]. With growing evidence of accelerated Greenland ice melt [Velicogna, 2009; Bamber *et al.*, 2012; Box and Colgan, 2013], increasing attention has been given to the role of meltwater in the high-latitude climate [Hu *et al.*, 2011; Weijer *et al.*, 2012; Rahmstorf *et al.*, 2015]. Proshutinsky *et al.* [2015] conjecture that the recent observed disruption of quasi-decadal oscillations of the Arctic wind-driven circulation regimes may be attributed to a surplus freshwater flux from Greenland. The underlying assumption is that a substantial volume of Greenland freshwater spreads laterally from the Greenland coast into convective sites of the interior sub-Arctic seas. On the other hand, observational studies suggest that freshwater predominantly flows in boundary currents along the margins of the Nordic Seas (the East Greenland Current) and the Labrador Sea (the West Greenland Current and the Labrador Current) [e.g., Bacon *et al.*, 2002; Dickson *et al.*, 2007; Sutherland and Pickart, 2008; Myers *et al.*, 2009]. Hence, most of the Greenland freshwater discharge should travel along the margins of the sub-Arctic seas on its way to the North Atlantic (Figure 1). It remains unclear how, where, and on what timescales this freshwater can impact convective regions [Moore *et al.*, 2015]. The rate of Greenland freshwater flux into the interior seas,

which is unknown, is also important. Previous model experiments have shown that the thermohaline circulation in the North Atlantic responds to freshwater perturbations on order of 0.1 Sv [Rahmstorf, 1995; Fanning and Weaver, 1997; Clark et al., 2002].

In order to understand the mechanisms of climate change driven by changes in AMOC in response to increased ice sheet melting, a number of “hosing” experiments have been performed with Ocean General Circulation Models and climate models [Huybrechts et al., 2002; Fichefet et al., 2003; Ridley et al., 2005; Jungclauss et al., 2006; Gerdes et al., 2006; Stouffer et al., 2006; Swingedouw et al., 2007; Vizcaino et al., 2008; Hu et al., 2011]. Overall, these experiments show different AMOC states as a function of freshwater flux into the North Atlantic. These numerical experiments are not designed to address any of the questions raised above. Typically in these hosing experiments surplus freshwater flux is imposed over some region of the North Atlantic. Although such simplification of freshwater input into the North Atlantic seems to have low impact on climate simulations [Kleinen et al., 2009], misrepresentation of actual pathways of Greenland freshwater into the ocean may be important for the realistic responses of thermohaline processes in the sub-Arctic seas, as discussed in Stammer [2008], Marsh et al. [2010], and Weijer et al. [2012].

In order to investigate the influence of surplus Greenland Ice Sheet meltwater on the sub-Arctic seas and North Atlantic several high-resolution ocean experiments have been conducted. A model study of Marsh et al. [2010] includes 0.25° and 1° NEMO simulations integrated for 8 years. They found that the freshwater signal tends to stay along the narrow boundary current of the Labrador Sea and only a very small volume of freshwater has reached the interior Subpolar Gyre in the simulation after 8 years. The largest volume of surplus freshwater accumulated in Baffin Bay. The authors mention that the East Greenland Coastal Current, which acts as a freshwater conduit, is not resolved in the NEMO

experiments. *Bacon et al.* [2014] conclude that at least 1/12 degree model resolution is needed to resolve freshwater fluxes associated with East Greenland Coastal Current.

Another recent model study of *Weijer et al.* [2012] compares the adjustment of the AMOC to the surplus of Greenland freshwater in multidecadal simulations with 0.1° (“strongly eddying”) and 1° (“non-eddying”) global configurations from the Los Alamos Parallel Ocean Program (POP). In these experiments, a passive tracer is released along with the freshwater anomaly around Greenland in order to diagnose the freshwater propagation rate. The numerical experiments predict rapid spreading of the tracer at shallow depths over the sub-Arctic seas (within a year) with a distinct delay (about a year) in the arrival time for the tracer in the interior Nordic Seas. Both POP configurations predict decline in convective activity in both the Labrador Sea and Nordic Seas. The strongest freshening is simulated in the Labrador Sea and Baffin Bay, whereas the decline in surface salinity in the Nordic Seas is much weaker.

The goal of this paper is to investigate the pathways, mechanisms and time scales of Greenland meltwater propagation within the sub-Arctic seas with particular focus on the Nordic Seas, the Labrador Sea, and Baffin Bay (Figure 1). We start our analysis with a description of the Greenland Ice Sheet runoff data used in the model experiments (section 2) and a brief overview of observed salinity changes in the sub-Arctic seas that could provide evidence of the influence of surplus Greenland freshwater influence (section 3). Design and analysis of numerical experiments employing three regional coupled ocean–sea ice models with different resolution forced by realistic Greenland freshwater fluxes are discussed in section 4. The experiments use a passive tracer released at exact locations of the Greenland freshwater sources. The model results are analyzed from the perspective of observed salinity changes in the sub-Arctic seas (section 5).

2. Greenland Freshwater Flux

2.1. Data

Greenland freshwater fluxes, between 1958 and 2010, from *Bamber et al.* [2012] are employed to investigate the fate of this freshwater and assess its role in thermohaline variability over the past decade. The data are a monthly gridded product (5x5km grid) with realistic geographic distribution and temporal variability (Figure 2a). Greenland runoff was derived from a reconstruction of the surface mass balance of the Greenland Ice Sheet and surrounding tundra using a high-resolution regional climate model, RACMO2 [*Ettema et al.*, 2009] forced with ERA-40 re-analysis data. The runoff from the ice sheet and surrounding tundra was combined with observations of solid ice discharge derived from satellite observations of ice velocity to produce the total freshwater flux.

2.2. Variability and Trends, 1990-2010

According to *Bamber et al.* [2012], total Greenland freshwater flux has been predominantly $>1000 \text{ km}^3 \text{ yr}^{-1}$ since 1998 having increased from 870—900 $\text{km}^3 \text{ yr}^{-1}$ in the early 1990s to 1100—1200 $\text{km}^3 \text{ yr}^{-1}$ in the late 2000s (Figure 2b). A somewhat smaller estimate of the total Greenland freshwater flux is reported in *Box and Colgan* [2013], but it did not include tundra runoff that accounts for an additional 100-200 $\text{km}^3 \text{ yr}^{-1}$. However, their estimate however indicates a similar increase of the Greenland freshwater flux of around 200 $\text{km}^3 \text{ yr}^{-1}$ between 1990 and 2010.

The volume of freshwater supplied by the Greenland Ice Sheet [*Bamber et al.*, 2012] is roughly 29% - 42% of the total annual river runoff into the Arctic Ocean (3500 $\text{km}^3 \text{ yr}^{-1}$ according to *Curry and Mauritzen* [2005] or 2500 $\text{km}^3 \text{ yr}^{-1}$ as reported by *Aagaard and Carmack* [1989] based on climatological data of the 1980s). The change of the freshwater flux between 1990 and 2010 is equivalent to a freshwater input increase of 6.3 mSv to 10.5

mSv. This increase is smaller than those of an earlier study of *Gregory and Lowe* [2000] noting a 17.4 mSv increase. In addition, between 2004 and 2009 mass loss from the Canadian Arctic Archipelago (CAA), primarily into Baffin Bay, changed from 31 to 92 km³ yr⁻¹ [*Gardner et al.*, 2011], an increase from about 1 to 3 mSv.

Estimated from the Greenland freshwater data of *Bamber et al.* [2012], the cumulative Greenland freshwater flux anomaly between 1990-2010 is 2830±337 km³ relative to the pre-1990 mean flux of 876 km³ yr⁻¹. This amount is 28% of the freshwater volume advected to the sub-Arctic from the Arctic Ocean during the 1970s GSA event, totaling 10,000 km³ [*Dickson et al.*, 1988]. If the Greenland freshwater flux increases at the same rate for the next 50 years, the surplus freshwater will reach the GSA freshwater volume by the mid 2060s. It is noteworthy that rates of mass loss from Greenland have been steadily increasing since 2010 [*Helm et al.*, 2014].

3. Observed Salinity Changes in the Sub-Arctic Seas

Recent changes in salinity fields in the sub-Arctic seas are discussed below in an attempt to identify observational evidence of freshwater release from Greenland that has been increasing since the early 1990s.

3.1. Baffin Bay

Hydrographic characteristics of Baffin Bay are largely determined by outflow from the Arctic Ocean and inflow from the Labrador Sea that enters the bay as the West Greenland Current (WGC, Figure 1). Greenland freshwater flux to Baffin Bay has substantially increased since the early 1990s (200—250 km³ yr⁻¹) contributing about 316 km³ yr⁻¹ in the late 2000s (Figure 2b). This is ~13% of the freshwater transport through the northern channels of the Canadian Arctic Archipelago (CAA) estimated as 2450 km³ yr⁻¹ combining estimates in *Peterson et al.* [2012], *Münchow and Melling* [2008], *Rabe et al.* [2012], *Agnew et al.* [2008], and *Kwok* [2005].

A detailed analysis of warming and freshening in Baffin Bay over 1916-2003 based on historical hydrographic data was conducted by *Zweng and Munchow* [2006]. The majority of the observations used were collected between 1950 and the early 2000s. The study revealed significant freshening over the shelf and slope regions of Baffin Bay in the layer between 50–200 m depth range. The largest freshening (-0.086 ± 0.039 decade⁻¹) was found on the Baffin Island shelf. A smaller freshening (from -0.048 to -0.066 decade⁻¹) was reported for the continental slopes off Baffin Island. The surface waters of the Greenland shelf and slope regions to the north of Davis Strait had similar freshening trends (-0.04 decade⁻¹). The authors suggest Greenland meltwater runoff might have contributed to this freshening. Recently *Myers and Ribergaard* [2013] also found freshening in the upper layer of Disko Bay, as well as offshore in the West Greenland Current in the 2000s as compared to the 1980s and 1990s.

Moored observations in Davis Strait described in *Curry et al.* [2014] indicate slight freshening of several water mass classes (Figures 3a-c). Although no statistical significance can be drawn from these short time series, all but two water masses (West Greenland Irminger Water, WGIW and Transitional Water, TrW) tend to have negative salinity anomalies in recent years relative to the 2003-2013 mean (Figures 3b and c). UK Met Office Hadley Centre objectively mapped monthly salinity data (version EN.4.1.1) [*Good et al.*, 2013] have been examined in the region (Figure 3d). Salinity anomalies relative to the 1960-1990 monthly climatology indicate apparent freshening signal in the upper 50 m over 1990 to 2014 in BB. Over the past decade, fresh surface water has expanded southwards (by about 10° latitude) into the northern Labrador Sea.

3.2. The Labrador Sea

The Labrador Sea receives freshwater exported from the Arctic Ocean through the CAA via Davis Strait and Hudson Strait and through Fram Strait with the EGC and WGC (Figure 1). Additionally, the Labrador Sea also receives salty modified Atlantic Water originating from the North Atlantic Current (called the Subpolar Mode Water in [McCartney and Talley, 1982]). The warm and salty Atlantic water flows along the continental slope around the basin as the Irminger Current. It is generally accepted that eddies shed by the Irminger Current control the heat and salt budget of the interior Labrador Sea [e.g., Straneo, 2006], governing winter convective activity when the intermediate and deep waters of the North Atlantic are formed [Lazier et al., 2002]. Deep convection controls formation and properties of Labrador Sea Water that occupies the central Labrador Sea [Talley and McCartney, 1982]. Deep winter convection in 1987—1994 produced an anomalously high volume of the deepest and densest Labrador Sea Water [Yashayaev, 2007]. After 1994, the freshening of the water column reversed, especially in the layers below 1000 m attributed to weaker convection in the central Labrador Sea with exception to the deep convection events during winter of 2007-2008 [Yashayaev and Loder, 2009].

Hydrographic observations in the central Labrador Sea indicate considerable variability of the temperature and salinity characteristics of the Labrador Sea Water over the last four decades [Yashayaev, 2007]. Oceanographic monitoring of the Labrador Sea was mostly done on the basis of annual hydrographic sections across the central Labrador Sea (for more details, see Yashayaev [2007]) until 2002 when the international Argo program (<http://www.argo.ucsd.edu>) was launched. In the early 1970s, the Labrador Sea had a noticeable freshening trend that continued until 1995 [Dickson et al., 2002; Zweng and Munchow, 2006; Yashayaev and Seidov, 2015]. During 1988 through 1994, strong freshening was observed in the upper 1500 m in the Labrador Sea caused by the formation of a large volume of cold and fresh Labrador Sea Water [Sy et al., 1997; Lazier et al., 2002]. After

1995, the salinity trend changed to positive and remained positive at least through the late 2000s. *Yashayaev and Loder* [2009] analyzed time evolution of hydrography fields in the central Labrador Sea on the basis of the Argo float data and in situ observations. Their results demonstrated a strong positive salinity anomaly developed in the 200-500 m layer during 2000-2009 with the highest salinity >34.9 in 2008 (compared to 34.78-34.82 in the late 1980s). Recent Argo observations in the Labrador Sea show a slight decrease in salinity in the 200-800 m layer after 2012 caused by several deep convection events, with the strongest convection occurred in 2014 [*Kieke and Yashayaev*, 2015]. Freshening in the upper 200m has been observed in the western Subpolar Gyre region and the Labrador Sea since 2010 [*Beszczyńska-Moller and Dye*, 2013; *Yashayaev et al.*, 2015].

3.3. The Nordic Seas

Hydrographic characteristics of the upper ocean water masses in the Nordic Seas are influenced by the northward flowing Atlantic Water and southward flowing Polar Water from the Arctic Ocean carried by the EGC [*Aagaard et al.*, 1985]. Whereas the eastern and western parts of the Nordic seas are largely determined by the characteristics of the Polar Water and Atlantic Water, respectively, the interior part, that includes convective regions in the Iceland and Greenland seas, has a more complex relationship between hydrography and contributions from Polar Water and Atlantic Water. The Greenland and Iceland seas have many features in common [*Malmberg and Jonsson*, 1997]. Hydrography of the upper layer in both seas is determined by water mass transformations and modifications through lateral advection of Polar Water and Atlantic Water from the rims of the seas and convection driven by surface cooling and salt flux during ice formation. The North Atlantic water undergoes substantial cooling and freshening as it mixes with the ambient water before entering the Greenland

Gyre. Upon mixing, the two water masses form the upper Greenland and Iceland water mass, sometimes called Arctic Surface Water [*Carmack, 1990*].

Large salinity changes have been observed in the Nordic Seas since the 1950s. A long-term freshening of the North Atlantic in surface, intermediate and deep water masses has been reported in *Curry et al. [2003]*, *Dickson et al. [2003]*, and *Curry and Mauritzen [2005]*. Hydrographic observations conducted in the Greenland and Iceland seas (regions “GS” and “IS” in Figure 1) reveal strong negative salinity anomalies (<34.3 to 34.5) in the upper 25-m layer during the late 1960s-1970s, late 1980s, and the late 1990s (Figure 4). The anomalies are related to the 1970s, 1980s and 1990s GSAs, respectively [*Dickson et al., 1988; Belkin et al., 1998; Belkin, 2004*]. In the mid 2000s, high salinity (>34.7) was observed in both seas in the layer from 0 to 200m. The positive salinity event was replaced by a strong freshening signal (<34.5) in 2008 in the upper Greenland Sea (above 50 m) but not in the Iceland Sea (Figure 4). A persistent salinity increase in the layers below 50–75 m was observed in the regions, especially apparent in the Greenland Sea. The observations in Figure 4 agree with the previous studies that reported a positive salinity trend in the Nordic Seas since the late 1990s [*Avsic et al., 2006; Hátún et al., 2005; Holliday et al., 2008; Somavilla et al., 2013*]. Recent analysis of Argo float observations in the Nordic Seas, between 2001-2007, shows a slightly positive salinity trend for the upper 500 m [*Latarius and Quadfasel, 2010*]. Increased salinity in the upper Nordic Seas is related to increased salinity in the Atlantic water flowing into the sub-Arctic seas [*Walczowski and Piechura, 2006; Holliday et al., 2008; Yashayaev and Seidov, 2015*], likely related to increased salinity in subtropical and mid-latitudes Atlantic Water [*Hátún et al., 2005*].

To summarize, historic observations in Baffin Bay show persistent freshening over the last two decades, whereas salinity in the other sub-Arctic seas has both positive and negative anomalies. Thus, no definite conclusion can be made on the linkage between

observed salinity anomalies in the sub-Arctic seas and increased Greenland freshwater discharge observed during last decades. We believe that more information and additional studies are needed to reveal mechanisms and processes regulating freshwater fluxes and fresh water transformations under different complex processes. A better knowledge about freshwater pathways, accumulation rates and time scales of freshwater propagation is necessary in order to understand the disagreement between proposed freshening of the sub-Arctic seas caused by surplus Greenland freshwater flux and the observed salinity changes. Note that observational studies referenced in this section were primarily focused on hydrographic changes in the water column below the near-surface layer (upper 50–200 m). However, results from model experiments with passive tracers described in the following sections suggest that Greenland freshening should primarily manifest in the in the upper 200 m and mostly in the 0–50 m layer.

4. Results from the Passive Tracer Experiments

Three groups participating in the Forum for Arctic Ocean Modeling and Observational Synthesis (FAMOS) project have run ocean-sea ice models with passive tracers released at the Greenland freshwater sources. The models used in this study are from the Florida State University (AO-HYCOM), the University of Alberta (NEMO-LIM2), and the Institute of Computational Mathematics and Mathematical Geophysics (ICMMG). The simulations have different model forcing and configurations. The length of integration is 14 years. The models have different resolutions; AO-HYCOM has a high-resolution (0.08°) configuration and NEMO-LIM2 and ICMMG have 0.25° and 0.5° grid spacing, respectively. The only coordinated forcing is the Greenland freshwater flux data (described in section 2.1). In the model experiments, Greenland freshwater is tracked by a passive tracer released at the locations of Greenland freshwater sources (Figure 2a). The tracer is treated as a scalar

variable and its time evolution is described by scalar advection and diffusion equations similar to temperature and salinity. Specifics of the models and experiments are summarized in Appendix 1.

4.1. Evolution of Tracer Concentration in the Upper Ocean

Time evolution of the passive tracer concentration in the simulations (Figures 5—7) reveals general agreement among the models. In particular, during the first several years, the models simulate rapid spreading of the freshwater tracer into the Labrador Sea (mainly with the Labrador Current) and Baffin Bay where the highest concentration is maintained throughout the simulation. On the eastern side of Greenland, the tracer is exported from the Nordic Seas by the EGC into the Labrador Sea and the Subpolar Gyre region, and then the tracer is advected back into the Nordic Seas with the North Atlantic current. Yet there are obvious discrepancies in the simulated fields. The differences among the models are found in tracer concentration (and mass) in the sub-Arctic seas, tracer pathways and spreading into the interior basins, and timing of tracer advection into the Nordic Seas.

Specifically, during the first 5 years of the AO-HYCOM simulation the tracer stays within the EGC. There is an indication of some advection of the tracer with the East Icelandic Current to the east (Years 1, 2, 5, and 6). However, the interior Nordic Seas show no signature of the tracer until year 6. After that, the interior Nordic Seas slowly fill up with the tracer being brought by the North Atlantic Current after the tracer has travelled around the Subpolar Gyre (Figure 5).

The NEMO-LIM2 simulation (Figure 6) agrees with AO-HYCOM showing very limited lateral advection of the tracer off the EGC during the first 5 years of the simulation. Yet in contrast to AO-HYCOM, tracer spreading into the interior Nordic Seas by means of the west-to-east Jan Mayen current is evident in NEMO-LIM2, . The timing of tracer

propagation into the convective regions of the Nordic Seas is remarkably similar between the two models. Nevertheless, there is a notable distinction in how this propagation is simulated. In NEMO-LIM2 the tracer spreads from the south with the North Atlantic Current and from west off the EGC, whereas in AO-HYCOM the tracer is mainly advected by the North Atlantic Current.

The ICMMG simulation (Figure 7) qualitatively agrees well with both AO-HYCOM and NEMO-LIM2. Yet the ICMMG simulation does not predict any substantial tracer spreading off the EGC into the interior Nordic Seas, unlike the NEMO-LIM2 simulation. In ICMMG, it takes much longer (10 years) for the tracer to travel around the Subpolar Gyre before returning to the Nordic Seas with the North Atlantic Current, compared to AO-HYCOM and NEMO-LIM2. Another discrepancy between ICMMG and the two other models is sluggish propagation of the tracer into the interior of the basins in the ICMMG simulation. For example, propagation of the tracer in Baffin Bay is predominantly along the coast in the ICMMG simulation. By contrast, there is substantial lateral advection of the tracer into the interior Baffin Bay in both AO-HYCOM and NEMO-LIM2. Similarly in the Labrador Sea and the Nordic Seas, ICMMG simulates a slower rate of tracer spreading into the interior basins.

Another disagreement between ICMMG and the two other models is the evolution of the tracer concentration to the north and west off Greenland. ICMMG simulates an extensive spread of the tracer into the Arctic Ocean from the northern Greenland coast, whereas in both AO-HYCOM and NEMO-LIM2 simulations tracer propagation into the Arctic Ocean is limited and confined to the narrow shelf north of Greenland and the CAA.

4.2. Tracer budget analysis

In order to characterize propagation and accumulation of the tracer in the sub-Arctic seas, a tracer budget is calculated for the six regions shown in Figure 1. The study area is divided into three basins: the Nordic Seas (NS), the Labrador Sea (LS), and Baffin Bay (BB). Within the NS and LS, convective sites in the Greenland Sea (GS), the Iceland Sea (IS), and interior Labrador Sea (IL) are designated. For each region, passive tracer storage (volume-integrated tracer content or mass, M_{tr}) is calculated as

$$M_{tr}(t) = \int \int \int_V C_{tr}(x, y, z, t) dV, \quad (1)$$

where C_{tr} is tracer concentration (kg m^{-3}). Within a region, tracer fluxes (F_{tr}) are estimated for each segment bounding the region by integrating over the total water depth (H) along the segment l of length L

$$F_{tr} = \int_L \int_H C_{tr}(l, z, t) \mathbf{u}(l, z, t) \cdot \mathbf{n}(l) dz dl \quad (2)$$

where \mathbf{n} is an inward unit normal vector (positive flux into the region).

4.2.1. The Nordic Seas

All the models show a gradual increase of the volume-integrated tracer content signifying tracer accumulation in the Nordic Seas region over the simulation time (Figure 8a). However, the models disagree on the evolution rate of the tracer content. In the AO-HYCOM simulation, concentration of the tracer starts growing after the first 3 years. In the other two simulations, the content demonstrates a nearly steady increase over the whole integration time. All models show a negative flux across Denmark Strait (section 1) owing to the tracer export with the EGC (Figures 8 b—d); all models simulate a reasonably robust seasonal signal in this transport. In AO-HYCOM (Figure 8b), the tracer net flux is negative during the

first 5 years when tracers are exported out of the region. After 5 years of the simulation, tracer inflow to the Nordic Seas starts increasing through Iceland-Faroe-Shetland segments (sections 2 and 3 corresponding to the green and blue lines, respectively) and the net oceanic flux becomes positive exceeding the outflow through section 1. In NEMO-LIM2 (Figure 8c) the net flux is negative dominated by the EGC export of the tracer during the simulation. Contrary to the AO-HYCOM results, tracer fluxes across Iceland-Faroe-Shetland segments exhibit only minor increase after year 6. The increase in tracer content in the NS is due to the tracer advection with the East Icelandic and Jan Mayen currents in NEMO-LIM2 (Figure 6). Differing from AO-HYCOM, ICMMG (Figure 8d) simulates considerably smaller tracer inflow into NS with the North Atlantic Current through sections 2 and 3. The inflow cannot compensate tracer outflow through Denmark Strait. The imbalance in the net advective tracer fluxes is compensated by internal tracer flux along the Greenland coast, whereas in the AO-HYCOM the net advective flux is positive after 5 years indicating that tracer accumulation in the NS region is mainly attributed to the tracer advection by the North Atlantic Current.

4.2.2. The Greenland Sea

Tracer accumulation in the GS deduced from the time series of the volume-integrated tracer content is largest for the AO-HYCOM model (Figure 9a). During the first 2-3 years, there is no tracer signature in the region for all models. After several years, the tracers start spreading into the interior Greenland Sea. However, the timing and the propagation rate differ among the models. In AO-HYCOM, the tracer content starts growing after 4 years. During the last 4 years of the simulation, the accumulation rate slows down. Both NEMO-LIM2 and ICMMG simulate accumulation of the tracers in the GS at a slower rate than in AO-HYCOM.

There is good agreement between the general trend of tracer fluxes in AO-HYCOM and NEMO-LIM2, with much smaller fluxes in general for the ICMMG simulations (Figures 9b and d). Tracer fluxes across sections 9 and 12 dominate but nearly cancel each other due to the throughflow of the recirculating Atlantic waters in the northern side of the GS. It is the balance between the fluxes through the eastern and southern sections that determines the tracer budget inside the box. In the AO-HYCOM experiment, the inflow across the eastern section (10) is mostly positive and the southern flux oscillates between positive and negative. The fluxes in ICMMG remain near zero through year 13.

Tracer fluxes across sections 9 and 12 in NEMO-LIM2 have a strong seasonal signal due to the propagation path of the tracer from the EGC (Figure 6), which has strong seasonal change in the tracer concentration related to seasonal variability of Greenland freshwater flux (Figure 2a). After year 8, the seasonal signals of the fluxes across sections 9 and 12 are superimposed on positive and negative trends, respectively, attributed to tracer that has travelled around the Subpolar Gyre and has been advected into the GS with the Norwegian Atlantic Current.

4.2.3. The Iceland Sea

Accumulation of the passive tracer in the IS is delayed by about 1 year in NEMO-LIM2, 5 years in AO-HYCOM and 10 years in ICMMG (Figure 10a). The earlier accumulation of the tracer in the IS region in NEMO-LIM2 is related to eastward tracer transport with the East Icelandic Current (Figure 6) that is not evident in the other two simulations.

Tracer fluxes across the sections have noticeable disagreement among the models (Figures 10b—d). AO-HYCOM simulates major tracer outflow through the southern section (16) and the inflow is mainly through the northern (14) and eastern (15) boundaries. Tracer

flux through the western boundary (13) is intermittent but becomes predominantly positive by the end of the simulation. Note that after year 6, tracer flux across the southern section (16) is always negative, whereas the northern flux (14) remains positive. Similarly, ICMMG simulates persistent positive tracer flux through the northern section and negative tracer flux through the southern section (Figure 10d), however the northern inflow dominates the outflow (note the different scales). In the NEMO-LIM2 simulation, the fluxes through sections 14 and 16 are more sporadic and do not demonstrate any persistence, in contrast to AO-HYCOM and ICMMG. The oscillations across sections 13 and 16 are related to variability of the East Icelandic Current and the Jan Mayen Current that are substantial tracer sources for the interior Nordic Seas in the model.

4.2.4. The Labrador Sea

All model experiments predict an overall increase of the passive tracer content in the LS during the simulation (Figure 11a). The rate of tracer increase is similar in all simulations. Tracer fluxes across the northern (7) and southern (8) sections exhibit strong seasonality associated with seasonal changes of meltwater flux from Greenland coast (Figures 11b—d). In AO-HYCOM and NEMO-LIM2, the southern flux through section 8 is positive during the first year of the simulation, and then it becomes predominantly negative. The shift is due to the fact that during the first year, the tracer is imported into the region with the WGC. After the tracer propagates around the Labrador Sea, it is exported out to the North Atlantic with the Labrador Current. Tracer flux between the Labrador Sea and Baffin Bay (section 7) oscillates on the seasonal time scale remaining mainly positive during the first half of the year and is reversed during the second half. Such a strong seasonal cycle is consistent with hydrographic observations in the West Greenland Current [Myers *et al.*, 2009; Curry *et al.*, 2014; Rykova *et al.*, 2015]. In the ICMMG simulation, the tracer flux across section 8

remains predominantly positive through year 10 caused by tracer influx from the North Atlantic. In contrast to the AO-HYCOM and NEMO-LIM2, the tracer is exported to Baffin Bay (through section 7) with little tracer export from the bay until year 11.

4.2.5. The Interior Labrador Sea

The rate and timing of tracer accumulation in the IL region agree between AO-HYCOM and ICMMG, whereas NEMO-LIM2 predicts a near-linear increase of the tracer until year 12 when tracer mass reaches maximum that is more than 2 times higher than in the AO-HYCOM simulation and nearly 5 times higher than in ICMMG (Figure 12a). Tracer fluxes across the sections bounding the IL box indicate different tracer advection pathways in the model experiments (Figures 12b—d). In AO-HYCOM, the fluxes do not have any persistent pattern. On average, tracer content in the IL is determined by the influx through the eastern (section 18) and northern (17) sides and the outflows through the western (20) and southern (19) sides. The NEMO-LIM2 experiment has the strongest tracer inflow through the southern section (19), while tracer flux across the northern section (17) is primarily negative. Similar to AO-HYCOM, in ICMMG the tracer budget in the IL box is largely influenced by the flux across the eastern side (18). Tracer outflow from the IL mostly occurs through the northern section (17).

4.2.6. Baffin Bay

The BB region shows accumulation of passive tracer in all experiments with the accumulation rate differing among the models (Figure 13a). In AO-HYCOM, the tracer content rapidly increases during the first 2 years. After that, the tracer content slowly increases with superimposed seasonal variability. The NEMO-LIM2 simulation predicts faster and more persistent tracer accumulation resulting in two time more tracer mass by the

end of the integration compared to AO-HYCOM. Results from the ICMG experiment demonstrate a near-linear increase in the tracer content in the BB region through the simulation also exceeding twice the tracer content in the AO-HYCOM simulation by the end of the experiment. Fluxes across the northern (6) and the southern (7) sections are dominated by seasonal change (Figures 13b—d). During the first half of the year the tracer is advected from the Nares Strait but during the second half of the year the tracer is advected from the Labrador Sea through Davis Strait. The seasonality is attributed to wind climatology in the area with strong southerly winds in winter and weaker northerly winds in the eastern bay during summer [Tang *et al.*, 2004].

4.3. Tracer Distribution in the Labrador Sea, the Nordic Seas and Baffin Bay

The numerical experiments illustrate that the tracer has been redistributed by ocean circulation within the entire sub-Arctic region over a relatively short time. At the beginning of the experiment, the tracer is predominantly located in the Labrador Sea and Baffin Bay (Figures 5—7). As the tracer spreads over the domain, the tracer content increases in the Nordic Seas. Analysis of the volume-integrated tracer content (section 4.2) reveals that tracer mass increases in all regions. Part of this increase is due to the overall increase of tracer mass in the domain and comparatively minor leakage of the tracer to the southern North Atlantic and the Arctic Ocean. However redistribution of the tracer within the sub-Arctic seas driven by the ocean circulation influences the rate of tracer accumulation rate in the basins.

In order to compare the accumulation rates, the ratio of volume-integrated tracer content (mass) in the Labrador Sea, the Nordic Seas and Baffin Bay relative to the total tracer mass released is analyzed (Figure 14). This ratio in the Labrador Sea and Baffin Bay decreases with time in all experiments. By the end of the simulations, the ratio of tracer mass in the Labrador Sea and Baffin Bay relative to the total tracer mass released is highest in the

ICMMG simulation compared to the other two models. This ratio indicates that the interior Labrador Sea (Figure 14, green numbers) loses the tracer over the course of the AO-HYCOM and ICMMG simulations, while in the NEMO-LIM2 simulation there is an accumulation of tracer in the region. In the Nordic Seas, the evolution of the tracer mass ratio is in the opposite sense between the AO-HYCOM the other two models indicating substantial differences in the simulated tracer spreading in the region. All the simulations indicate accumulation of tracer inside the GS and IS boxes (Figure 14, blue and red numbers). By the end of the simulations, the three basins contain 29.5% (AO-HYCOM), 28% (NEMO-LIM2), and 38.2% (ICMMG) of the total tracer mass. The remainder of the tracer is accumulated in the Subpolar Gyre region and in the Arctic Ocean. It is of note that the largest fraction of the tracer mass remains in the sub-Arctic seas (including the Subpolar Gyre region) after 14 years of the simulations, demonstrating that the tracer has been accumulating in the region while being transported around the sub-Arctic seas and propagating into the interior regions. The result corroborates previous studies that discuss recirculation of negative and positive salinity anomalies within the Labrador – Subpolar Gyre – Nordic Seas [Belkin, 2004; Yashayaev and Seidov, 2015].

4.4. Tracer spreading into the deep layers

Observations suggest that water masses propagate predominantly along isopycnal surfaces [e.g., Ledwell *et al.*, 1993]. Greenland freshwater spreads into the sub-Arctic seas remaining near the surface and mixing with ambient water. In the sub-Arctic seas, freshwater is mixed downward by haline and thermal convection [e.g., Watson *et al.*, 1999; Rudels *et al.*, 2012]. Thus, Greenland meltwater ultimately penetrates into the deep layers in the convective areas. Several studies suggest that freshwater can be mixed downward into the deep layers and then laterally advected to distant basins. In the numerical experiments, the tracers being

released in the near-surface layer tagging buoyant meltwater quickly penetrate the subsurface layers through mechanical mixing and deep convection. Spurious diapycnal mixing can also enhance vertical propagation of the tracer in the non-isopycnal NEMO-LIM2 and ICMMG models [Bleck, 2002; Hill *et al.*, 2012].

The models predict tracer maximum concentration in the near-surface layer (see sections in Figure 15). However, the maximum is strongly diffused in NEMO-LIM2 and ICMMG occupying depths between 200 and 500 m, by contrast, it remains within a narrow layer (upper 100 m) in AO-HYCOM. In the AO-HYCOM experiment, the deepest propagation of the tracer occurs in the North Atlantic (Figures 15a and b). In the Nordic Seas, the tracer is at notably shallower depths (~800m) with distinct minima in the Greenland Sea (shallower than ~700 m). Similarly, in Baffin Bay the tracer has not spread deeper than ~800m. Development of deep local tracer maxima is evident in the southern Labrador Sea by the end of year 7 (also observed in Figure 14a). After 14 years of the model experiment, the passive tracer has penetrated depths exceeding 1600-1800 m in the North Atlantic including the Labrador Sea and the southern Irminger Sea. The concentration decreases with depth and is maximal in the surface layer (<200 m).

In contrast to AO-HYCOM, NEMO-LIM2 simulates tracer spreading through the whole water column in Baffin Bay (Figures 15c and d). In the Labrador Sea, the tracer is distributed through most of the water column down to ~2800 m after 7 years and all the way to the bottom after 14 years. This is related to excessive Labrador Sea convection in this simulation. After 7 years, the tracer propagation shallows downstream of the North Atlantic flow with very shallow tracer signatures in the Nordic Seas owing to insufficient propagation of the tracer into this region. By the end of the simulation, the tracer spreads to the bottom in the North Atlantic and below 600 m in the Nordic Seas.

In the ICMMG experiment (Figures 15e and f), vertical propagation of the tracer generally agrees with AO-HYCOM. After 14 years, the model simulates deeper propagation of the tracer in the Labrador Sea and the North Atlantic basin and substantially shallower propagation in the Nordic Seas and Baffin Bay. ICMMG tracer distribution after 7 years differs from both the AO-HYCOM and NEMO-LIM2 simulations. Specifically, in ICMMG the tracer spreads deeper in the Labrador Sea, compared to the AO-HYCOM but shallower than in NEMO-LIM2. There is no tracer signature in the Nordic Seas in the ICMMG. There is good agreement in tracer depth in Baffin Bay between AO-HYCOM and ICMMG. After 14 years, tracer distribution in the North Atlantic basin in the ICMMG looks more different from AO-HYCOM. Interestingly there is no tracer in the near-surface layers in the North Atlantic basin in the ICMMG simulation, suggesting the advection origin of the subsurface tracer presence in this part of the section. The absence of the tracer in the central Labrador Sea and Nordic Seas is due to the tracer propagation with the boundary currents in this simulation.

5. Discussion

5.1. Pathways of Greenland Freshwater

5.1.1. Horizontal propagation

The pathways of Greenland freshwater propagation follow the general circulation in the sub-Arctic seas (Figure 1). The presented tracer experiments demonstrate relatively rapid spreading of the passive tracer in the sub-Arctic seas reaching the most remote regions within 5 – 7 years. Estimated travel time of the tracer in the model experiments agrees well with the estimates of propagation rates of the GSAs in the sub-Arctic inferred from observations [Dickson *et al.*, 1988; Belkin *et al.*, 1998; Belkin, 2004; Yashayaev and Seidov, 2015]. Based on the observed cooling and freshening signals in the region, Dickson *et al.* [1988] traced the

spreading of the 1970s GSA and suggested that it took the GSA 7-8 years to propagate from Fylla Bank (off the southwestern Greenland coast) to the Norwegian Sea and 9-10 years to Spitsbergen. Upstream, propagation timescales for the GSA were 1 year longer from the eastern Greenland coast (north of Iceland). The GSAs of the 1980s and the 1990s spread faster propagating from Fylla Bank to the Norwegian Sea in about 5-6 years [Belkin *et al.*, 1998; Belkin, 2004]. The major inflow of the tracer into the Nordic Seas in AO-HYCOM and ICMMG occurs with the North Atlantic Current. This result is in agreement with other studies where the dominant role of negative salinity anomalies carried by the Atlantic inflow on the freshening signal in the Nordic Seas has been proposed [e.g., Glessmer *et al.*, 2014; Reverdin, 2014].

The discrepancies in the numerical solutions are apparent in tracer propagation into the interior regions of the sub-Arctic seas. There are markedly stronger horizontal gradients in tracer concentration fields simulated in ICMMG, compared to the other two models. In the coarse-resolution ICMMG simulation, the interior slowly fills with the tracer that tends to remain with the current following the boundary regions (Figure 7). For example, in ICMMG the tracer is advected into the Nordic Seas after 5 years of the simulation in agreement with the AO-HYCOM experiment, yet it takes another 7 – 8 years for the tracer to spread into the interior GS and IS in the ICMMG simulation compared to 2 – 3 years in the AO-HYCOM experiment (Figure 5). One possible reason for the observed discrepancies in the tracer distribution between ICMMG and the other two models could be the coarse grid spacing of the former. The coarse resolution of the ICMMG model does not permit the simulation of mesoscale eddies that may be a key mechanism for advection of tracer into the interior regions.

This assumption is supported by previous studies suggesting a leading role of small-scale eddies in lateral advection and spreading of both freshwater and Atlantic Water in the

627 Labrador Sea [Saenko *et al.*, 2014] and the Nordic Seas [Budeus and Ronski, 2009;
 628 Yashayaev and Seidov, 2015]. For example, Budeus and Ronski [2009] reported a patchy
 629 distribution of Atlantic Water observed inside the Greenland Gyre. Diameters of these
 630 “Atlantic Water patches” were only 20 km with vertical extent varying from 200 to 1000 m.
 631 The findings of Budeus and Ronski [2009] support the idea of eddy transport of Atlantic
 632 Water into the interior Greenland Gyre (see also Yashayaev and Seidov [2015] for an
 633 interesting analogy of Atlantic Water circulation and a pinball machine). This view is distinct
 634 from the idea of a diffusion type of penetration that assumes gradual lateral spreading
 635 resulting in a smooth and steady transition from salinity and temperature values at the
 636 boundaries of the gyres to the interior values [e.g., Karstensen *et al.*, 2005]. This type of
 637 advection is present in the ICMMG (non-eddy) simulation.

638 The above discussion raises the question about the meaning of “eddy-resolving” with
 639 respect to Arctic Ocean models. An “eddy-resolving” model has to be capable to adequately
 640 representing the eddy field, i.e. model horizontal grid spacing should be at least two grid
 641 points per Rossby radius of deformation and the grid spacing is measured as the grid-diagonal
 642 distance (so called “effective spacing” in Hallberg [2013]). In the Nordic Seas, the first
 643 baroclinic Rossby radius of deformation is $\sim 4\text{--}8$ km [Nurser and Bacon, 2013]. Hence, an
 644 “eddy-resolving” model should have a grid spacing about 2–4 km. Only AO-HYCOM with
 645 0.08° horizontal grid (effective spacing $\sim 4\text{--}5$ km in the Greenland Sea) is close to this
 646 estimate still being marginal between “eddy-resolving” and “eddy-permitting” (1 grid point
 647 per radius [Nurser and Bacon, 2013]). No other model from the previous studies mentioned
 648 earlier (section 1) satisfies this resolution criterion. With >20 km horizontal grid spacing, in
 649 the ICMMG model tracer propagation into the gyres is dominated by horizontal diffusion,
 650 which is much slower than transport and mixing by eddies in the other two models.

The apparent disagreement in the NEMO-LIM2 solution with the other models is that NEMO-LIM2 shows stronger tracer propagation from the EGC with the Jan Mayen Current. The causes of this intensification are not clear. We speculate that this may be related to misrepresentation of the narrow East Greenland Coastal Current, which serves as a freshwater conduit in the Nordic Seas. This problem has been discussed with respect to another 0.25° NEMO simulation in *Marsh et al.* [2010]. Alternatively, it could be due to the stronger East Icelandic Current in NEMO-LIM2 compared to the other two models.

5.1.2. Non-Eddying and Eddy Fluxes

The general circulation in the sub-Arctic seas follows the boundaries of the basins suggesting no tracer propagation into the interior Labrador and Nordic Seas. The AO-HYCOM and NEMO-LIM2 simulations, however, indicated tracer spreading into the interior regions (section 4) in striking contrast to the coarse-resolution ICMMG simulation that showed very minor propagation of the tracer into the interior boxes. Visual inspection of figures 5–7 support the rationale suggested above that different eddy activity explains the different spreading of the tracer into the interior regions. To validate this assumption, eddy tracer flux and mean (non-eddy) tracer flux are calculated by separating velocities and concentration into time-mean velocity ($\bar{\mathbf{u}}$) and tracer concentration (\bar{C}_{tr}) and corresponding time-fluctuating components or eddy terms (\mathbf{u}' and C'_{tr}). After time averaging, the tracer fluxes may be separated into mean and eddy fluxes

$$\overline{\mathbf{u} \cdot C_{tr}} = \bar{\mathbf{u}} \cdot \bar{C}_{tr} + \overline{\mathbf{u}' \cdot C'_{tr}}, \quad (3)$$

where the overbar denotes 30-day time averaging.

Figure 16 presents one-year-averaged mean and eddy tracer fluxes for year 9 from the simulations. The mean tracer flux (Figures 16 a, c, and e) highlights the pathways of the tracer in the sub-Arctic seas that follow the general large-scale circulation. The general

features of the mean tracer flux are similar in the models. Yet there are notable discrepancies in the smaller-scale pathways. For example, in contrast to the higher-resolution simulations, in ICMMG boundary currents are broad occupying nearly whole Baffin Bay. At the same time, the interior Nordic Seas is nearly quiescent. Note that the mean tracer flux in the Nares Strait is northward, in contradiction to AO-HYCOM and NEMO-LIM2 (as well as the observations). The eddy tracer fluxes (Figures 16 b, d, and f) are predominantly directed normal to the mean flux, indicating lateral advection of the tracer into the interior regions. All the models simulate increased eddy tracer flux along southwestern and southeastern Greenland coast. The magnitude of the eddy tracer flux in the Labrador Sea is markedly higher in AO-HYCOM compared to the other two experiments (note the natural-log scale). Also AO-HYCOM fields predict a local maximum of the eddy tracer fluxes in the Norwegian Sea that is smaller in NEMO-LIM2 and is absent in ICMMG.

Next, time-integrated eddy tracer flux (F_{eddy}) and time-mean tracer flux (F_{mean}) in the upper 150 m are calculated for the interior boxes in the Greenland, Iceland, and Labrador seas

$$F_{mean}(t) = \int_t \int_H \int_L \bar{u}_\perp \cdot \bar{C}_{tr} dl dz dt, \quad (4)$$

$$F_{eddy}(t) = \int_t \int_H \int_L \overline{u'_\perp \cdot C'_{tr}} dl dz dt, \quad (5)$$

where u_\perp is the velocity component normal to the section.

Shown in Figure 17 are time-integrated eddy tracer flux (Eq. 4) and mean tracer flux (Eq. 5) calculated for years 8—12. All the models demonstrate modest contribution of eddy tracer flux compared to the mean flux in the Nordic Seas. This is an expected result from the previous analysis of Figure 16. Strikingly different from the Nordic Seas is the substantial contribution of eddy flux in the Labrador Sea in the AO-HYCOM simulation. The amount of tracer advected by eddies into the IL box is higher than the amount advected by the mean

fluxes. This result is different from the NEMO-LIM2 and ICMMG simulations where the eddy tracer flux is negligibly small in the IL region.

Thus, analysis of the tracer fluxes simulated in the models has demonstrated the predominant role of the mean tracer flux into the interior Nordic Seas. The eddy tracer flux dominates the non-eddy tracer flux in the interior Labrador Sea in the AO-HYCOM simulation. Calculated eddy fluxes do not support the idea of eddy advection as the dominant mechanism of tracer transport into the interior Nordic Seas, in contrast to the Labrador Sea.

5.2. Greenland Freshwater Influence on Salinity in the sub-Arctic Seas from AO-HYCOM

The simulations with the passive tracer have demonstrated rapid tracer propagation over the sub-Arctic seas suggesting widespread influence of Greenland freshwater flux in the region in agreement with the hypothesis discussed in Section 1. The tracer concentration increases in all the sub-Arctic seas during 14 years of model simulations indicating accumulation of Greenland freshwater in the basins. The results imply the growth of freshwater content that should manifest in increasing negative salinity anomaly in the upper sub-Arctic seas. Nevertheless, there is no obvious observational evidence that could relate freshening signals in the region with Greenland freshwater flux. According to the model results, accumulation over the last two decades of surplus Greenland freshwater should have resulted in a negative salinity trend in the near-surface layer. Instead, observed salinity changes alternate between positive and negative anomalies with no persistent temporal pattern (as discussed in section 3). In the following section, the influence of the Greenland freshwater flux on salinity changes in the sub-Arctic region is analyzed using results from only the AO-HYCOM simulation, which has the highest resolution.

5.2.1. Greenland Freshwater Flux and Salinity Change in the Sub-Arctic

The AO-HYCOM experiment was initialized from another simulation that had been integrated without Greenland runoff. In the tracer experiment, Greenland freshwater flux is “turned on” at the first time step of the model integration. Hence, there is a discrete jump in the freshwater forcing from the initial conditions with zero Greenland runoff to $>1000 \text{ km}^3 \cdot \text{yr}^{-1}$ (0.032 Sv) in the simulation. This provides easier detection and tracking of Greenland freshwater propagation and associated freshwater anomalies in the sub-Arctic seas. The drawback of the experiment design is that simulated negative salinity changes in the sub-Arctic seas can be too extreme for direct comparison with observational records. A better approach would be to initialize the experiment with the ocean fields from another experiment forced by realistic Greenland runoff, although in practice this still complicates the process of detection of freshwater anomalies in the basin. The AO-HYCOM results are employed here to establish relationships between Greenland freshwater flux and salinity changes in the sub-Arctic. This relationship will be used to estimate freshening in the sub-Arctic seas caused by the surplus Greenland freshwater flux in 1990-2010.

Both freshwater and salinity changes in the sub-Arctic basins have linear trends during the simulation (Figure 18). For every region (Figure 1), salinity is spatially averaged within the slab 0-400 m inside the 500-m isobath. A linear relationship is sought between the spatially averaged salinity and cumulative Greenland freshwater mass Q_r

$$Q_r(t) = \int_0^t F_r dt, \quad (6)$$

where F_r is Greenland freshwater flux. The tracer experiments have demonstrated a delay between the time when freshwater (or passive tracer) is released at the Greenland coast and the time of appearance of the freshwater signature in the sub-Arctic basins (Figure 4). The time lag (τ) varies for the basins from 1 year for Baffin Bay to more than 5 years for the Greenland and Iceland Seas. Thus, the following regression is fit to the data

$$dS(t) = \alpha_0 + \alpha_1 \cdot Q_r(t - \tau), \quad (7)$$

where dS is salinity change averaged in the upper 400 m within a sub-Arctic basin at time t (Figure 18b). The time lag is defined iteratively until the best fit to the data is provided, based on the highest value of the coefficient of determination (R^2). The high values of R^2 in the analyzed cases indicate a strong linear relationship between the Greenland freshwater flux and the magnitude of salinity anomaly in the regions.

The next step is to “remap” dS onto the observational Greenland freshwater changes reported in *Bamber et al.* [2012]. Greenland freshwater flux increased by $\sim 200 \text{ km}^3 \text{ yr}^{-1}$ ($\sim 0.006 \text{ Sv}$) from 1990 to 2010. Assuming a linear increase in the Greenland freshwater flux over this period, Eq. (7) provides the estimated salinity change (dS) in the six sub-Arctic regions for 1990–2015 (Figure 18c). The magnitude of salinity anomaly is modest in the basins, especially in the interior Nordic Seas (GS and IS), suggesting a relatively small contribution of the Greenland runoff on the freshening of the sub-Arctic seas over 20 years of increased Greenland freshwater flux. According to Figure 18c, by 2010 salinity anomaly should be -0.05 in the GS, -0.07 in the IS, and -0.08 in the IL as a result of surplus Greenland freshwater flux.

An important caveat here is that salinity changes of the North Atlantic Current have not been taken into account. According to long-term observations in the Iceland-Faroe-Shetland section and in the southwestern Nordic Seas salinity of Atlantic water increased by $0.15 - 0.2$ in the 2000s compared to the long-term mean [*Holliday et al.*, 2008; *Yashayaev and Seidov*, 2015]. Such a strong positive salinity anomaly would dominate and undermine the impact of Greenland freshwater runoff on salinity in the sub-Arctic seas.

6. Summary

As Greenland Ice Sheet melting has been accelerating, increased freshwater discharge into the ocean can have dramatic consequences for thermohaline circulation of the sub-Arctic seas. Generally accepted ramifications of the surplus freshwater flux are increased water column stability and weakening of deep convection in the interior Labrador and Nordic Seas, not to mention sea level rise as the most prominent consequence. Several hypotheses of the current and future climate changes in the Arctic and North Atlantic have been suggested on the basis of this assumption. Yet pathways and time scales of freshwater propagation in the sub-Arctic seas are not known. The influence of Greenland freshwater on convective regions remains elusive. Moreover, recent observational records demonstrate increasing salinity in the upper Nordic Seas and the Labrador Sea during the 2000s contradicting the anticipated freshening caused by accelerated Greenland Ice Sheet melt. Although freshening in the 0–200 m layer in the Labrador Sea was observed during the 2010s [Beszczynska-Möller and Dye, 2013; Yashayaev *et al.* 2015]. In order to address these uncertainties, three numerical experiments were conceived during FAMOS discussions. In these experiments employing AO-HYCOM, NEMO-LIM2, and ICCMG ocean-sea ice models, a passive tracer was continuously released at Greenland freshwater source sites to track propagation of the freshwater in the sub-Arctic seas.

Results from the tracer experiments demonstrate general agreement among the models on timing and propagation of the tracer in the sub-Arctic seas. The tracer follows the large-scale ocean circulation pattern with the EGC, WGC and the Subpolar Gyre from Greenland to the North Atlantic and with the North Atlantic Current to the Nordic Seas. The tracer quickly propagates into Baffin Bay and the Labrador Sea. The interior Labrador Sea is impacted by the tracer within the first 2 years of the simulations. The major pathway of the

tracer to the Nordic Seas is via the North Atlantic Current after it has traveled around the Subpolar Gyre. It takes from 3 (AO-HYCOM) to 5 (NEMO-LIM2 and ICMMG) years for the tracer to reach the south Nordic Seas via the North Atlantic Current and another 2 years to enter convective sites in the interior Nordic Seas (except for ICMMG where tracer spreading into the interior region was not substantial over the simulation time). The Iceland Sea receives the tracer via the Iceland Current shortly after the simulation begins, yet the amount of this influx is not substantial, except in the NEMO-LIM2 simulation. Positive trends in the time series of the tracer mass inside the basins indicate that a steady state has not been reached in the simulation and the tracer was still accumulating in the sub-Arctic seas by the end of the model runs. Nevertheless, the accumulation has substantially slowed in Baffin Bay and the Labrador Sea to the end of the experiments.

The model experiments predict highest tracer concentration in Baffin Bay with the second highest concentration in the Labrador Sea. The concentration in these basins quickly reached a quasi-steady state (during the first 4-5 years). In the rest of the domain, the concentration increased as the tracer propagated with the Subpolar Gyre and the North Atlantic Current towards the Nordic Seas. During the 14-year time interval, the tracer accumulated in the sub-Arctic seas. However, the simulations were not sufficiently long to reach a steady state in tracer concentration within the domain precluding determination of the residence time of the tracer in the sub-Arctic basins.

The noticeable discrepancy in the numerical solutions is tracer propagation into the convective regions in the Labrador Sea and the Nordic Seas. In the ICMMG simulation the interior regions are filled slowly with the tracer resulting in strong horizontal gradients of the tracer concentration, whereas the other two models simulate faster propagation of the tracer into the interior seas. The disagreement in the simulations is probably due to inability of the coarse-resolution ICMMG to represent tracer advection by small-scale eddies that may play

an important role in distributing water from the currents following the margins of the sub-Arctic seas. Calculated eddy tracer fluxes demonstrated substantially higher eddy activity in the AO-HYCOM simulation, contrasting NEMO-LIM2 and ICMMG. Nevertheless, calculated eddy tracer flux into the Greenland and Iceland interior boxes did not demonstrate noticeable contribution of eddies to the total tracer flux. By contrast, eddy flux was a substantial contribution to the tracer flux into the interior Labrador Sea.

Modeled vertical distribution of tracer in the water column has a near-surface (top 200 m) maximum in the sub-Arctic seas. Simulated vertical penetration of the tracer is deepest in the North Atlantic where the tracer spreads down to the near-bottom layers in AO-HYCOM and NEMO-LIM2 and down to ~2000m in ICMMG. The models disagree on the vertical tracer distribution in Baffin Bay where AO-HYCOM and ICMMG predicts tracer penetration down to ~800m and NEMO-LIM2 mixes the tracer all the way to the bottom. The causes of this disagreement are unclear and need further investigation.

Accumulation of the tracer in the interior regions of the sub-Arctic seas supports the idea of Greenland meltwater influence of the thermohaline processes and convection in the region. These results, however, are not directly supported by observations. No persistent negative salinity trends can be found in the reported hydrographic changes in the sub-Arctic region except for Baffin Bay. The estimates of the impact of the surplus Greenland freshwater flux on salinity changes in the upper 400 m layer suggest noticeable yet not dramatic freshening (from -0.04 to -0.12) of the sub-Arctic seas, especially in the interior regions. However, the magnitude of this freshening signal is smaller than the observed salinity increase in the Atlantic Water by 0.15–0.2 during the 2000s [Holliday *et al.*, 2008]. The salinity increase in the Atlantic inflow can be linked to the propagating positive salinity anomaly from the subtropical Atlantic Ocean [Hátún *et al.*, 2005; Straneo and Heimbach, 2013]. This positive salinity anomaly counteracts the freshening signal caused by Greenland

freshwater. The results of the model experiments also suggest that the accumulation of Greenland freshwater in the sub-Arctic seas caused by continuing Greenland Ice Sheet melt can amplify freshening in the Nordic Seas and the Labrador Sea during the period when salinity of the Atlantic inflow from southern North Atlantic decreases.

While the results here attest to efficient transport of freshwater derived from Greenland melt into the sub-Arctic seas, feedbacks and implications of this freshwater, as well as its relative importance with respect to influxes from the Arctic Ocean and Atlantic, remain an open question.

7. Acknowledgements

Davis Strait mooring observational data used in this study are available for download at <http://iop.apl.washington.edu/data.html>, with the timeseries also available through ACADIS, the repository for US Arctic Observing Network data. The study was part of U.S. National Science Foundation Freshwater Initiative (2004-2007) and the International Polar Year and Arctic Observing Network (2007-2010) programs under grants OPP0230381 and OPP0632231. Additional support was provided by the Department of Fisheries and Oceans, Canada. Hydrographic measurements (salinity, temperature, and pressure from bottle and CTD data) in the Greenland and Iceland seas from 1950 to 2010 have been used in this study are freely available from the ICES Dataset on Ocean Hydrography (www.ocean.ices.dk) and the PANGAEA database (www.pangaea.de). Greenland freshwater flux data analyzed in this study is that presented in *Bamber et al.* [2012] and is available on request as a gridded product. The analysis of salinity in Baffin Bay and the northern Labrador Sea has been performed on the basis of the UK Met Office Hadley Center observations dataset version EN.4.1.1 [*Good et al.*, 2013] (downloaded from <http://www.metoffice.gov.uk/hadobs/en4/> on

874 17 November, 2015). D. Dukhovskoy was funded by NASA JPL IOVWST and ONR NOPP.
875 E. Chassignet was supported by the ONR. HYCOM simulation was supported by a grant of
876 computer time from the DoD High Performance Computing Modernization Program at NRL
877 SSC. P.G. Myers and X. Hu were supported by NSERC grants RGPIN 227438-09, RGPIN
878 04357 and RGPCC 433898. G. Platov was supported by RFBR grants 13-05-00480, 14-05-
879 00730, and 15-05-02457. We thank W. Weijer (LANL) and I.M. Belkin (URI) for their
880 thoughtful suggestions and comments.
881

Appendix A. Characteristics of the numerical models

A1. AO-HYCOM

The high resolution 0.08° regional Arctic Ocean HYbrid Coordinate Ocean Model (HYCOM) coupled to the Los Alamos National Laboratory Community Ice Code (CICE) [Hunke and Lipscomb, 2008] is used in this application (hereinafter referenced as AO-HYCOM). HYCOM is a primitive equation generalized coordinate (hybrid) ocean model [Bleck, 2002; Chassignet *et al.*, 2006]. The model domain is a subset of the global HYCOM [Metzger *et al.*, 2014] north of $\sim 38^\circ\text{N}$. The computational grid of the AO-HYCOM is a Mercator projection from the southern boundary to 47°N . North of 47°N , it employs an orthogonal curvilinear Arctic dipole grid [Murray, 1996]. The model employs 32 hybrid vertical coordinate layers with potential density referenced to 2000m and includes the effect of thermobaricity [Chassignet *et al.*, 2003]. The model is forced with atmospheric fields (2-m air temperature, 2-m atmospheric humidity, surface shortwave and longwave heat fluxes, and precipitation) that are derived from hourly fields of the Climate Forecast System Reanalysis (CFSR) [Saha *et al.*, 2010]. Wind stress is estimated from the Cross-Calibrated Multi-Platform Ocean Surface Wind vector Analyses (CCMP) [Atlas *et al.*, 2011]. The atmospheric fields employed for this experiment cover the period from 2004-2010 and have been recycled to provide forcing for 14 years. Surface latent and sensible heat fluxes, along with evaporation, are calculated using bulk formulas during model run time. The bulk transfer coefficients are parameterized following Kara *et al.*'s [2000] algorithm.

The model is initialized from an existing HYCOM-CICE simulation that was run with no Greenland runoff, and is integrated continuously for 14 years. Lateral open boundaries are derived from a climatology from the 0.08° Global HYCOM hindcast [Metzger *et al.*, 2014]. Thus, there is no interannual variability in the oceanic forcing superimposed at the open boundaries. It is noteworthy that surface salinity relaxation is minimal in the simulation with

the e-folding relaxation scale of $1.59\text{e-}8 \text{ s}^{-1}$ corresponding to the restoring time scale of 4 year⁻¹.

Greenland freshwater sources are incorporated into AO-HYCOM using monthly inter-annual gridded data (Fig. 1b). The amount of tracer is proportional to the local freshwater flux rate and thus, tracer input replicates seasonal and interannual variability of the Greenland freshwater discharge. In HYCOM, runoff from a single source is distributed over several grid cells. From the beginning of the experiment, the passive tracer is continuously released in the upper 6m at every model grid cell that has non-zero Greenland freshwater influx along the Greenland coast. The tracer is prescribed in the model by relaxing tracer concentration in the specified locations, which are ocean grid cells nearest to the freshwater sources along the Greenland coast, to the maximum concentration value that is defined as follows. For the given Greenland freshwater flux at some location along the coast (F_r , m³/s) tracer concentration (kg/m³) in the grid cell is defined as

$$C_{tr} = \frac{F_r \cdot t_{rlx} \cdot \rho_{tr}}{\Delta z \cdot \Delta x \cdot \Delta y}, \quad (\text{A1})$$

where F_r is Greenland runoff flux at a given location (m³ s⁻¹), ρ_{tr} is tracer density (1000 kg m⁻³), t_{rlx} is tracer relaxation time scale (1 day, here) and Δz , Δx and Δy are layer thickness and horizontal grid spacing (m). Tracers are released in the upper 2 layers in the grid cells where Greenland runoff is prescribed.

A.2. NEMO-LIM2

The Nucleus for European Modelling of the Ocean model (NEMO v3.4 – Madec, 2008) is used in this experiment. The Arctic Northern Hemisphere Atlantic configuration (ANHA4) was based on the 0.25° tripolar grid extracted from the NEMO ORCA025 configuration developed within the Mercator-Ocean and DRAKKAR collaboration [Barnier *et al.*, 2007]. The model consists of 50 vertical levels with a 1 m top layer decreasing in

931 resolution with increasing depth. The ANHA4 domain is contained within open boundaries at
932 20°S latitude and Bering Strait. Lateral boundaries are free slip. Lateral mixing varies
933 horizontally according to a bi-Laplacian operator with a horizontal eddy viscosity of $1.5 \times$
934 $10^{11} \text{ m}^4 \text{ s}^{-1}$. For tracer lateral diffusion the model uses an isopycnal Laplacian operator with a
935 horizontal eddy diffusivity of $300 \text{ m}^2 \text{ s}^{-1}$. Vertical mixing at sub-grid scales was
936 parameterized using a turbulent kinetic energy (TKE) closure model [Madec *et al.*, 1998;
937 Axell, 2002]. Background vertical eddy viscosity and diffusivity are $10^{-4} \text{ m}^2 \text{ s}^{-1}$ and $10^{-5} \text{ m}^2 \text{ s}^{-1}$
938 respectively.

939 The sea ice module is from the Louvain-la-Neuve sea-ice model (LIM2) [Fichefet and
940 Maqueda, 1997] with a modified elastic-viscous-plastic (EVP) ice rheology [Hunke and
941 Dukowicz, 1997]. No-slip and free-slip boundary conditions are applied for sea ice and ocean,
942 respectively.

943 The simulations presented here were forced with inter-annual atmospheric data
944 derived from the Canadian Meteorological Centre's Global Deterministic Prediction System
945 (CGRF, [Smith *et al.*, 2013]) with an hourly resolution in time and a spatial resolution of
946 0.45° longitude and 0.3° latitude (minimal spacing is $\sim 33 \text{ km}$ in the Labrador Sea).

947 The model is initialized with the output from the GLORYS2V3 reanalysis from
948 MERCATOR and then run from 2002-2010 and then recycled over 2004-2010 to provide a
949 14 year simulation. It is forced with monthly inter-annual runoff from Dai *et al.*, [2009].
950 Greenland freshwater sources are incorporated using monthly inter-annual gridded data of
951 [Bamber *et al.*, 2012]. Starting in January 2004, passive tracer is continuously released in the
952 upper 10 m of the model at every freshwater source on the Greenland coast, with 5 tracers
953 defined around the coasts of Greenland. The amount of tracer is proportional to the local
954 freshwater discharge rate and thus, tracer flux replicates seasonal and inter-annual variability

of the Greenland freshwater discharge. The local change of tracer concentration in the model grid cells is defined as

$$\Delta C_{tr} = \frac{F_r \Delta t}{\Delta z \Delta x \Delta y}. \quad (A2)$$

A.3. ICMMG

A regional model of the Arctic and Atlantic Oceans of the Institute of Computational Mathematics and Mathematical Geophysics (ICCMG) is configured from 20°S in the Atlantic Ocean to 60°N in the Pacific Ocean [Golubeva and Platov, 2007]. The horizontal computational grid is bipolar curvilinear and has an equatorial resolution of 0.5° (minimal spacing is ~19km in the study region). The model uses a hydrostatic primitive formulation of Navier-Stokes equations with the rigid-lid approximation. The sea ice model is version 3 of CICE [Hunke and Lipscomb, 2008].

The model is forced by wind stress, sensible and latent heat fluxes, precipitation and evaporation, solar and longwave radiation derived from the NCEP/NCAR reanalysis [Kanamitsu *et al.*, 2002]. Lateral open boundaries are provided by the PHC climatology (http://psc.apl.washington.edu/nonwp_projects/PHC/Climatology.html). The experiment is initialized from an existing 1948-2003 model run with no Greenland runoff. Greenland freshwater input is implemented in the model as surface mass and salt fluxes at the ocean grid cells closest to the freshwater sources on the Greenland coast. Similar to the other model experiments, the amount of tracer is proportional to the local freshwater discharge rate. Tracers are prescribed as mass flux ($\text{kg m}^{-2} \text{s}^{-1}$) into the near-surface layer calculated from the local runoff rate as

$$F_{tr} = \frac{F_r \rho_{tr}}{\Delta x \Delta y}. \quad (A3)$$

979
980
981

Table A1. Characteristics of the numerical models

	AO-HYCOM	NEMO-LIM2	ICMMG
Horizontal grid	Dipole curvilinear, 0.08°	Curvilinear, 0.25°	Dipole curvilinear, 0.5°
Vertical coordinates	32 hybrid layers	50 geopotential levels	38 geopotential levels
Free surface	Free surface, split time step	Linear filtered free surface ¹	Rigid-lid approximation
Baroclinic time step	240 s	1080 s	5400 s
Barotropic time step	7.5 s	none	Varying
Salinity relaxation	e-folding relaxation scale = $1.59 \times 10^{-8} \text{ s}^{-1}$.	No relaxation	No relaxation
Bathymetry	DBDBV2 ²	ETOPO1 ³ +GEBCOv1 ⁴	2.5km IBCAO ⁵
Scalar horizontal advection	Second-order flux-corrected transport	TVD ⁶	Ultimate QUICKEST ⁷
Horizontal diffusion	Laplacian diffusion = $u_d \times \Delta x$, u_d for scalars = 0.005 m/s, u_d for momentum = $2.86 \times 10^{-3} \text{ m/s}$. Biharmonic diffusion: $u_d \times \Delta x^3$ momentum dissipation = 0.03 m/s	Laplacian for tracer, maximum eddy diffusivity is $300 \text{ m}^2 \text{ s}^{-1}$ (proportional to grid size); Bilaplacian for momentum, maximum eddy viscosity is $-1.5 \times 10^4 \text{ m}^4/\text{s}$ (proportional to the cubic grid size)	Laplacian, $50 \text{ m}^2/\text{s}$
Vertical turbulence	KPP	TKE (turbulent kinetic energy) vertical mixing model	Richardson-based vertical mixing ⁸ and OPPS ⁹
Diapycnal diffusivity	$1 \times 10^{-7} / \text{buoyancy freq.}, \text{ m}^2 \text{ s}^{-1}$	None	None
Background diffusivity	$0.1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$	$0.1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$	$0.1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$
Background viscosity	$0.3 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$	$1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$	$1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$
Tracer input	Relaxed to tracer concentration based on local Greenland river runoff rate		Surface mass flux
Atmospheric forcing	CCMP winds and CFSR radiative fluxes	CGRF	NCEPR
Sea ice model	CICE v.4	LIM2	CICE v.3

982 ¹ Roullet and Madec (2000)
983 ² Digital Bathymetric Data Base Variable Resolution (DBDBV2)
984 ³ Global 1 min resolution relief dataset [*Amante and Eakins*, 2009]
985 ⁴ General Bathymetric Chart of the Oceans
986 ⁵ International Bathymetric Chart of the Arctic Ocean
987 ⁶ Total Variation Dissipation scheme [*Lévy et al*, 2001]
988 ⁷ [*Vested et al.*, 1992]
989 ⁸ [*Golubeva and Platov*, 2007]
990 ⁹ Ocean penetrative plume scheme [*Paluszkievicz and Romea*, 1997]
991

991

992 *References*

- 993 Aagaard, K. and E. C. Carmack (1989), The role of sea ice and other fresh-water in the Arctic
 994 circulation, *J. Geophys. Res.*, 94(C10), 14,485-14,498, doi: 10.1029/JC094iC10p14485.
- 995 Aagaard, K., J.H. Swift, and E.C. Carmack (1985), Thermohaline circulation in the Arctic
 996 Mediterranean Seas, *JGR*, 90(C3), 4833-4846.
- 997 Atlas, R., R. N. Hoffman, J. Ardizzone, S. M. Leidner, J. C. Jusem, D. K. Smith, and D.
 998 Gombos (2011), A cross-calibrated, multiplatform ocean surface wind velocity product
 999 for meteorological and oceanographic applications, *Bull. Amer. Meteor. Soc.*, 92, 157-
 1000 174. doi: 10.1175/2010BAMS2946.1.
- 1001 Avsic, T., J. Karstensen, U. Send, and J. Fischer (2006), Interannual variability of newly
 1002 formed Labrador Sea Water from 1994 to 2005. *Geophysical Research Letters* 33,
 1003 L21S02, doi:10.1029/2006GL026913.
- 1004 Axell, L. B. (2002), Wind-driven internal waves and langmuir circulations in a numerical
 1005 ocean model of the southern baltic sea. *J. Geophys. Res.*, 107, doi
 1006 :10.1029/2001JC000922.
- 1007 Bacon, S., G. Reverdin, I.G. Rigor, and H.M. Snaith (2002), A freshwater jet on the east
 1008 Greenland Shelf, *J. Geophys. Res.*, 107(C7), 1–5, doi:10.1029/2001JC000935.
- 1009 Bacon, S., A. Marshall, N. P. Holliday, Y. Aksenov, and S. R. Dye (2014), Seasonal
 1010 variability of the East Greenland Coastal Current, *J. Geophys. Res. Oceans*, 119, 3967–
 1011 3987, doi:10.1002/2013JC009279.
- 1012 Bamber, J., M. van den Boreke, J. Ettema, and J. Lenaerts (2012), Recent large increase in
 1013 freshwater fluxes from Greenland into the North Atlantic, *GRL*, 39, L19501, doi:
 1014 10.1029/2012GL052552.
- 1015 Barnier, B., et al. (2006), Impact of partial steps and momentum advection schemes in a
 1016 global ocean circulation model at eddy-permitting resolution, *Ocean Dyn.*, 56(5), 543–
 1017 567, doi:10.1007/s10236.
- 1018 Batchelor, C.L. and J.A. Dowdeswell (2014), The physiography of High Arctic cross-shelf
 1019 troughs. *Quaternary Science Reviews* 92, 68-96. DOI: 10.1016/j.quascirev.2013.05.025.
- 1020 Belkin, I. M., S. Levitus, J. I. Antonov, and S.-A. Malmberg (1998), “Great Salinity
 1021 Anomalies” in the North Atlantic, *Prog. Oceanogr.*, 41, 1 – 68.
- 1022 Belkin, I. (2004), Propagation of the “Great Salinity Anomaly” of the 1990s around the
 1023 northern North Atlantic, *Geophys. Res. Lett.* 31, doi: 10.1029/2003GL019334.
- 1024 Beszczynska-Moller, A. and S.R. Dye (Eds.), *ICES Report on Ocean Climate 2012*. ICES
 1025 Cooperative Research Report No. 321, Copenhagen, Denmark, 73 pp.
- 1026 Bleck, R. (2002), An oceanic general circulation model framed in hybrid isopycnic Cartesian
 1027 coordinates. *Ocean Model.* 4, 55–88.
- 1028 Blindheim J. and S. Oesterhus (2005). The Nordic seas, main oceanographic features. In:
 1029 Drange H, Dokken T, Furevik T, Gerdes R, Berger W, editors. *The Nordic Seas: An*
 1030 *Integrated Perspective*. Washington, American Geophysical Union, p. 11-38.
- 1031 Bönisch G, J. Blindheim, J.L. Bullister, P. Schlosser, and D.W.R. Wallace (1997), Long-term
 1032 trends of temperature, salinity, density and transient tracers in the central Greenland Sea.
 1033 *J Geophys Res.*, 102, 553-71.
- 1034 Box, J.E. and W. Colgan (2013), Greenland ice sheet mass balance reconstruction. Part III:
 1035 Marine ice loss and total mass balance (1840-2010), *J. Climate*, 26, 6990-7002, doi:
 1036 10.1175/JCLI-D-12-00546.1.
- 1037 Budeus, G. and S. Ronski (2009), An integral view of the hydrographic development in the
 1038 Greenland Sea over a decade, *The Open Ocean Journal*, 3, 9-40.

- Carmack, E.C. (2000), Large-scale physical oceanography of polar oceans. In: The Freshwater Budget of the Arctic Ocean, ed. E.L. Lewis et al., Kluwer Academic Publishers, The Netherlands, 91-126.
- Carton, J.A., S. A. Grodsky, and H. Liu (2008), Variability of the Oceanic mixed layer, 1960-2004, *J. Climate* (21), 1029-1047.
- Castro de la Guardia, L., X. Hu, and P. G. Myers (2015), Potential positive feedback between Greenland Ice Sheet melt and Baffin Bay heat content on the west Greenland shelf, *Geophys. Res. Lett.*, 42, 4922–4930, doi:10.1002/2015GL064626.
- Clark, P.U., N.G. Pisias, T.F. Stocker, and A.J. Weaver (2002), The role of the thermohaline circulation in abrupt climate change, *Nature* 415, 863-869, doi:10.1038/415863a.
- Curry, B., C.M. Lee, B. Petrie, R.E. Moritz, and R. Kwok (2014), Multiyear Volume, Liquid Freshwater, and Sea Ice Transports through Davis Strait, 2004-10, *J. Phys. Oceanogr.* (44), 1244-1266.
- Curry, R.G. and C. Mauritzen (2005), Dilution of the northern North Atlantic Ocean in recent decades, *Science*, 308, 1772-1774.
- Curry, R.G., R. Dickson, and I. Yashayaev (2003), A change in the freshwater balance of the Atlantic Ocean over the past four decades, *Nature*, 426, 826-829.
- Chassignet, E.P., L.T. Smith, G.R. Halliwell, and R. Bleck. 2003. North Atlantic simulations with the HYbrid Coordinate Ocean Model (HYCOM): Impact of the vertical coordinate choice, reference pressure, and thermobaricity. *Journal of Physical Oceanography* 33:2,504–2,526, [http://dx.doi.org/10.1175/1520-0485\(2003\)033<2504:NASWTH>2.0.CO](http://dx.doi.org/10.1175/1520-0485(2003)033<2504:NASWTH>2.0.CO).
- Dai, A., T. Qian, K. E. Trenberth, and J. D. Milliman (2009), Changes in continental freshwater discharge from 1948 to 2004, *J. Clim.*, 22, 2773–2791
- Dickson, R.R., J. Meincke, S.-A. Malmberg, and A.J. Lee, (1988), The “Great Salinity Anomaly” in the Northern North Atlantic 1968–82, *Progress in Oceanography* 20, 103–151.
- Dickson, R.R., T.J. Osborn, W. Hurrell, J. Meincke, J. Blindheim, B. Adlandsvik, T. Vinje, G. Alekseev, W. Maslowski (2000), The Arctic Ocean response to the North Atlantic oscillation. *Journal of Climate* 13, 2671–2696.
- Dickson, R.R., Yashayaev, I., Meincke, J., Turrell, W., Dye, S., Holford, J., 2002. Rapid freshening of the deep North Atlantic over the past four decades. *Nature* 416, 832–837.
- Dukhovskoy, D.S., M.A. Johnson, and A. Proshutinsky (2004), Arctic decadal variability: An auto-oscillatory system of heat and fresh water exchange. *GRL*, 31, doi:10.1029/2—3GL019023.
- Fanning, A.F. and A.J. Weaver (1997), Temporal-geographical meltwater influences on the North Atlantic Conveyor: Implications for the Younger Dryas, *Paleoceanography*, 12(2), 307-320.
- Femke de Jong, M., H.M. van Aken, K. Vage, and R.S. Pickart (2012), Convective mixing in the central Irminger Sea: 2002-2010, *Deep-Sea Res. I*, 63, 36-51.
- Fichefet, T., and M. A. Morales Maqueda (1997), Sensitivity of a global sea ice model to the treatment of ice thermodynamics and dynamics, *Journal of Geophysical Research*, 102, 12,609–12,646.
- Fichefet, T., Poncin, C., Goosse, H., Huybrechts, P., Janssens, I., H. LeTreut (2003), Implications of changes in freshwater flux from the Greenland Ice Sheet for the climate of the 21st century. *Geophys. Res. Lett.* 30, 1911. doi:10.1029/2003GL017826.
- Fichefet, T., and M. A. Morales Maqueda (1997), Sensitivity of a global sea ice model to the treatment of ice thermodynamics and dynamics, *Journal of Geophysical Research*, 102, 12,609–12,646

- Gardner, A., G. Moholdt, B. Wouters, G.J. Wolken, D.O. Burgess, M.J. Sharp, J. G. Cogley, C. Braun, and C. Labine, (2011), Sharply increased mass loss from glaciers and ice caps in the Canadian Arctic Archipelago, *Nature*, 473, 357–360, doi:10.1038/nature10089.
- Gerdes, R., W. Hurlin, and S.M. Griffies (2006), Sensitivity of a global ocean model to increased run-off from Greenland. *Ocean Model.* 12, 416–435.
- Glessmer, M.S., T. Eldevik, K. Vage, J.E.O. Nilsen, and E. Behrens (2014), Atlantic origin of observed and modeled freshwater anomalies in the Nordic Seas, *Nature Geoscience*, 7, 801–805.
- Golubeva, E. N., and G. A. Platov (2007), On improving the simulation of Atlantic water circulation in the Arctic Ocean, *J. Geophys. Res.*, 112, C04S05, doi:10.1029/2006JC003734.
- Good, S. A., M. J. Martin and N. A. Rayner, 2013. EN4: quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *Journal of Geophysical Research: Oceans*, 118, 6704–6716, doi:10.1002/2013JC009067.
- Goosse, H., F. M. Selten, R. J. Haarsma, and J. D. Opsteegh (2002), A mechanism of decadal variability of the sea-ice volume in the Northern Hemisphere, *Clim.Dyn.*, 19, 61– 83, doi:10.1007/s00382-001-0209-5.
- Gregory, J.M. and J.A. Lowe (2000), Predictions of global and regional sea-level rise using AOGCMs with and without flux adjustment, *Geophysical Research Letters* 27, 3069–3072. doi:10.1029/1999GL011228.
- Griffies, S.M., R.C. Pacanowski, and R.W. Hallberg (2000), Spurious diapycnal mixing associated with advection in a z-coordinate ocean model, *Mon. Wea. Rev.* 128, 538–564.
- Haine, T.W.N., B. Curry, R. Gerdes, E. Hansend, M. Karcher, C. Lee, B. Rudels, G. Spreen, L. de Steur, K. D. Stewart, R. Woodgate (2015), Arctic freshwater export: Status, mechanisms, and prospects, *Global and Planetary Change* 125, 13–35.
- Hátún, H., A. B. Sando, H. Drange, B. Hansen, and H. Valdimarsson (2005), Influence of the Atlantic subpolar gyre on the thermohaline circulation, *Science*, 309, 1841– 1844.
- Helm, V., A. Humbert, and H. Miller (2014), Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2, *The Cryosphere*, 8(4), 1539–1559.
- Hill, C., D. Ferreira, J.-M. Campin, J. Marshall, R. Abernathey, N. Barrier (2012), Controlling spurious diapycnal mixing in eddy-resolving high-coordinate ocean models – Insights from virtual deliberate tracer release experiment, *Ocean Modelling*, 45–46, 14–26.
- Holliday, N. P., S.L. Hughes, S. Bacon, A. Beszczynska-Moller, B. Hansen, A. Lavin, H. Loeng, K.A. Mork, S. Osterhus, T. Sherwin, and W. Wlczowski (2008), Reversal of the 1960s to 1990s freshening trend in the northeast North Atlantic and Nordic Seas, *Geophys. Res. Lett.*, 35, L03614, doi:10.1029/2007GL032675.
- Hu, A., G.A. Meehl, W. Han, and J. Yin (2011), Effect of the potential melting of the Greenland Ice Sheet on the Meridional Overturning Circulation and global climate in the future. *Deep-Sea Research II*, doi:10.1016/j.dsr2.2010.10.069.
- Hunke, E.C. and J. K. Dukowicz (1997), An Elastic–Viscous–Plastic Model for Sea Ice Dynamics, *J. Phys. Oceanogr.*, 27, 1849–1867, doi: http://dx.doi.org/10.1175/1520-0485(1997)027<1849:AEVPMF>2.0.CO;
- Hunke, E.C., and W. Lipscomb (2008), CICE: The Los Alamos Sea Ice Model: Documentation and Software User’s Manual, Version 4.0, Technical Report LA-CC-06-012, Los Alamos National Laboratory, Los Alamos, NM.
- Huybrechts, P., et al., 2002. The response of the Greenland Ice Sheet to climate changes in the 21st century by interactive coupling of an AOGCM with a thermomechanical ice-sheet model. *Ann. Glaciol.* 35, 409–415.
- Ikeda, M. (1990), Decadal oscillations of the air-ice-ocean system in the northern hemisphere, *Atmos. Ocean*, 28, 106– 139.

- Ikeda, M., J. Wang, and J.-P. Zhao (2001), Hypersensitive decadal oscillations in the Arctic/subarctic climate, *Geophys. Res. Lett.*, 28, 1275–1278.
- Jahn, A. and M.M. Holland (2013), Implications of Arctic sea ice changes for North Atlantic deep convection and the meridional overturning circulation in CCSM4-CMIP5 simulations, *GRL*, 40(6), 1206–1211.
- Jungclaus, J.H., et al., 2006. Will Greenland melting halt the thermohaline circulation? *Geophys. Res. Lett.* 33, L17708. doi:10.1029/2006GL026815.
- Kanamitsu, M., W. Ebisuzaki, J. Woollen, S.-K. Yang, J. J. Hnilo, M. Fiorino, and G. L. Potter (2002), NCEP–DOE AMIP-II Reanalysis (R-2), *Bull. Amer. Meteor. Soc.*, 83, 1631–1643, doi: <http://dx.doi.org/10.1175/BAMS-83-11-1631>.
- Kara, A.B., Rochford, P.A., Hurlburt, H.E., 2000. Efficient and accurate bulk parameterizations of air-sea fluxes for use in general circulation models, *J. Atmos. Ocean Tech.*, 17, 1421–1438.
- Karstensen J, P. Schlosser, D.W.R. Wallace, J.L. Bullister, and J. Blindheim (2005), Water mass transformation in the Greenland Sea during the 1990s. *J Geophys Res C Oceans*, 110: doi:10.1029/ 2004JC002510.
- Kieke, D. and I. Yashayaev (2015), Studies of Labrador Sea Water formation and variability in the subpolar North Atlantic in the light of international partnership and collaboration. *Prog. Oceanogr.*, <http://dx.doi.org/10.1016/j.pocean.2014.12.010>.
- Kleinen, T., T. J. Osborn, and K. R. Briffa (2009), Sensitivity of climate response to variations in freshwater hosing location, *Ocean Dyn.*, 59, 509–521.
- Lazier, J., Hendry, R., Clarke, R., Yashayaev, I. & Rhines, P. (2002), Convection and restratification in the Labrador Sea, 1990–2000, *Deep Sea Research I* 49, 1819–1835.
- Ledwell, J.R., A.J. Watson, and C. Law (1993), Evidence for slow mixing across the pycnocline from an open-ocean tracer-release experiment. *Nature* 364, 701–703.
- Lévy, M., A. Estublier, and G. Madec (2001), Choice of an advection scheme for biogeochemical models, *Geophysical Research Letters*, 28 (19), 3725–3728.
- McCartney, M.S. and L.D. Talley (1982), The subpolar mode water of the North Atlantic Ocean. *J. Phys. Oceanogr.* 12(11), 1169–1188.
- Madec, G., P. Delecluse, M. Imbard, and C. Levy (1998), OPA 8 ocean general circulation model - reference manual. Tech. rep., LODYC/IPSL Note 11.
- Madec, G. (2008), NEMO ocean engine, Note du Pôle de modélisation, Institut Pierre-Simon Laplace (IPSL), France, No 27 ISSN No 1288-1619.
- Malmberg, S.-A. and S. Jonsson (1997), Timing of deep convection in the Greenland and Iceland seas, *ICES J. Marine Science*, 54, 300–309.
- Manabe, S. and R.J. Stouffer (1988), Two stable equilibria of a coupled ocean-atmosphere model, *J. Clim.*, 1, 841–866.
- Manabe, S. and R.J. Stouffer (1993), Century-scale effects of increased atmospheric CO₂ on the ocean-atmosphere system. *Nature* 364, 215–218.
- Manabe, S. and R.J. Stouffer (1995), Simulation of abrupt climate change induced by freshwater input to the North Atlantic ocean, *Nature*, 378, 165–167.
- Marsh, R., D. Desbruyères, J. L. Bamber, B. A. de Cuevas, A. C. Coward, and Y. Aksenov (2010), Short-term impacts of enhanced Greenland freshwater fluxes in an eddy-permitting ocean model, *Ocean Sci.*, 6, 749–760.
- Marsh, R., et al. (2015), NEMO-ICB (v1.0): interactive icebergs in the NEMO ocean model globally configured at eddy-permitting resolution, *Geoscientific Model Development*, 8(5), 1547–1562.
- Moore, G.W.K., K. Vage, R.S. Pickart, and I.A. Renfrew (2015), Decreasing intensity of open-ocean convection in the Greenland and Iceland seas, *Nature Climate Change*, doi: 10.1038.

1188 Münchow, A. and H. Melling (2008), Ocean current observations from Nares Strait to the
 1189 west of Greenland: Interannual to tidal variability and forcing. *J. Mar. Res.*, 66, 801–833,
 1190 doi:10.1357/002224008788064612.
 1191 Muenchow, A., K.K. Falkner, and H. Melling (2015), Baffin Island and West Greenland
 1192 Current Systems in northern Baffin Bay, *Progress in Oceanography* 132, 305–317
 1193 Murray, R.J. (1996), Explicit generation of orthogonal grids for ocean models. *Journal of*
 1194 *Computational Physics* 126:251–273, <http://dx.doi.org/10.1006/jcph.1996.0136>.
 1195 Myers, P. G., and M. H. Ribergaard (2013), Warming of the polar water layer in Disko Bay
 1196 and potential impact on Jakobshavn Isbræ, *J. Phys. Oceanogr.*, 43(12), 2629–2640,
 1197 doi:10.1175/JPO-D-12-051.1
 1198 Myers, P. G., C. Donnelly, and M. H. Ribergaard (2009), Structure and variability of the west
 1199 Greenland current in summer derived from 6 repeat standard sections, *Prog. Oceanogr.*,
 1200 80(1), 93–112, doi:10.1016/j.pocean.2008.12.003
 1201 Mysak, L. A., and S. A. Venegas (1998), Decadal climate oscillations in the Arctic: A new
 1202 feedback loop for atmospheric-ice-ocean interactions, *Geophys. Res. Lett.*, 25(19), 3607–
 1203 3610.
 1204 Nurser, A.J.G. and S. Bacon (2013). Eddy length scales and the Rossby radius in the Arctic
 1205 Ocean. *Ocean Science Discussions*, 10, 1807-1831.
 1206 Paluszkievicz, T. and R.D. Romea (1997), A one-dimensional model for the
 1207 parameterization of deep convection in the ocean. *Dynamics of Atmospheres and Oceans*
 1208 26, 95–130.
 1209 Peterson, I., J. Hamilton, S. Prinsenberg, and R. Pettipas (2012), Wind-forcing of volume
 1210 transport through Lancaster Sound, *J. Geophys. Res.*, 117, C11018,
 1211 doi:10.1029/2012JC008140.
 1212 Pickart, R.S., M.A. Spall, M.H. Ribergaard, G.W.K. Moore, and R.F. Milliff (2003), Deep
 1213 convection in the Irminger Sea forced by the Greenland tip jet, *Nature*, 424, 152-156.
 1214 Polyakov, I. V., et al. (2005), One more step toward a warmer Arctic, *Geophys. Res. Lett.*,
 1215 32, L17605, doi:10.1029/2005GL023740.
 1216 Pritchard, H.D., R.H. Arthern, D.G. Vaughan, and L.A. Edwards (2009), Extensive dynamic
 1217 thinning on the margins of the Greenland and Antarctic ice sheets, *Nature* 461, 971–975.
 1218 doi:10.1038/nature08471.
 1219 Proshutinsky, A., Bourke, R. H., and F. A. McLaughlin (2002), The role of the Beaufort Gyre
 1220 in Arctic climate variability: Seasonal to decadal climate scales. *Geophys. Res. Lett.*, 29,
 1221 (23).
 1222 Proshutinsky, A., R. Krishfield, M.-L. Timmermans, J. Toole, E. Carmack, F. McLaughlin,
 1223 W.J. Williams, S. Zimmermann, M. Itoh, and K. Shimada (2009), The Beaufort Gyre
 1224 Fresh Water Reservoir: State and variability from observations, *J. Geophys. Res.*,
 1225 doi:10.1029/2008JC005104.
 1226 Rabe, B., H. Johnson, A. Münchow, and H. Melling (2012), Geostrophic ocean currents and
 1227 freshwater fluxes across the Canadian Polar Shelf via Nares Strait. *J. Mar. Res.*, 70, 603–
 1228 640.
 1229 Rahmstorf, S. (1995), Bifurcations of the Atlantic thermohaline circulation in response to
 1230 changes in the hydrological cycle, *Nature*, 378, 145–149.
 1231 Rahmstorf, S. (2002), Ocean circulation and climate during the past 120,000 years, *Nature*,
 1232 419, 207–214.
 1233 Rahmstorf, S. (2003), Thermohaline circulation: The current climate. *Nature*, 421, 699.
 1234 Rahmstorf, S., J.E. Box, G. Feulner, M.E. Mann, A. Robinson, S. Rutherford, and E.J.
 1235 Schaernicht (2015), Exceptional twentieth-century slowdown in Atlantic Ocean
 1236 overturning circulation, *Nature Climate Change*, 5, 475–480, doi:10.1038/nclimate2554.

1237 Ridley, J.K., P. Huyberegts, J.M. Gregory, J.A. Lowe (2005), Elimination of the Greenland
 1238 Ice Sheet in a high CO₂ climate. *J. Clim.* 18, 3409–3427, doi: [http://dx.doi.org/](http://dx.doi.org/10.1175/JCLI3482.1)
 1239 10.1175/JCLI3482.1.
 1240 Rooth, C. (1982), Hydrology and ocean circulation. *Progr. Oceanogr.*, 11, 131-149.
 1241 Rudels, B. (2011), Volume and freshwater transports through the Canadian Arctic
 1242 Archipelago—Baffin Bay system, *J. Geophys. Res.*, 116, C00D10,
 1243 doi:10.1029/2011JC007019.
 1244 Rudels, B., M. Korhonen, G. Budeus, A. Beszczynska-Moller, U. Schauer, A. Nummelin, D.
 1245 Quadfasel, and H. Valdimarsson (2012), The East Greenland Current and its impacts on
 1246 the Nordic Seas: observed trends in the past decade, *ICES J. Marine Science*, 69(5), 841-
 1247 851, doi:10.1093/icesjms/fss079.
 1248 Rykova, T., F. Straneo, and A. S. Bower (2015), Seasonal and interannual variability of the
 1249 West Greenland Current System in the Labrador Sea in 1993–2008, *J. Geophys. Res.*
 1250 *Oceans*, 120, 1318–1332, doi:10.1002/2014JC010386.
 1251 Saha, S., S.S. Moorthi, H.-L. Pan, X. Wu, J. Wang, S. Nadiga, P. Tripp, R. Kistler, J.
 1252 Woollen, D. Behringer, H. Liu, D. Stokes, R. Grumbine, G. Gayno, J. Wang, yu-tai hou,
 1253 hui-ya chuang, hann-Ming h. juang, joe SeLa, Mark iredell, ruSS treadon, daryL kLeiSt,
 1254 PauL van deLSt, denniS keySer, john derber, Michael ek, jeSSe Meng, heLin Wei,
 1255 rongqian yang, StePhen Lord, huug van den dooL, arun kuMar, Wanqiu Wang, craig
 1256 Long, MuthuveL cheLLiah, yan Xue, boyin huang, jae-kyung ScheMM, WeSLey
 1257 ebiSuzaki, roger Lin, PingPing Xie, Mingyue chen, Shuntai zhou, Wayne higgins, cheng-
 1258 zhi zou, quanhua Liu, yong chen, yong han, Lidia cucuruLL, richard W. reynoLdS, gLenn
 1259 rutLedge, and Mitch goLdberg, (2010), The NCEP Climate Forecast System Reanalysis,
 1260 *Bull. Amer. Meteor. Soc.*, 91, 1015-1057, doi: 10.1175/2010Bams3001.1.
 1261 Smith, G. C., F. Roy, P. Mann, F. Dupont, B. Brasnett, J.-F. Lemieux, S. Laroche, and S.
 1262 Blair, 2014: A new atmospheric dataset for forcing ice–ocean models: Evaluation of
 1263 reforecasts using the Canadian global deterministic prediction system. *Quart. J. Roy.*
 1264 *Meteor. Soc.*, 140, 881–894, doi:10.1002/qj.2194.
 1265 Stammer, D. (2008), Response of the global ocean to Greenland and Antarctic ice melting,
 1266 *JOURNAL OF GEOPHYSICAL RESEARCH*, VOL. 113, C06022,
 1267 doi:10.1029/2006JC004079
 1268 Straneo, F. (2006), Heat and freshwater transport through the central Labrador Sea, *J. Phys.*
 1269 *Oceanogr.*, 36, 606–628.
 1270 Stommel, H. (1961), Thermohaline convection with two stable regimes of flow, *Tellus*, 2,
 1271 224–230.
 1272 Stouffer, R.J., J. Yin, J.M. Gregory, K.W. Dixon, M.J. Spelman, W. Hurlin, A.J. Weaver, M.
 1273 Eby, et al. (2006). Investigating the causes of the response of the thermohaline circulation
 1274 to past and future climate changes, *J. Climate*, 19, 1365-1387.
 1275 Straneo, F. and P. Heimbach (2013), North Atlantic warming and the retreat of Greenland's
 1276 outlet glaciers, *Nature*, 504, 36-43, doi: doi:10.1038/nature12854.
 1277 Sutherland, D. and R.S. Pickart (2008), The East Greenland Coastal Current: Structure,
 1278 variability, and forcing, *Prog. Oceanogr.*, 78, 58–77.
 1279 Swingedouw, D., P. Braconnot, and O. Marti (2006), Sensitivity of the Atlantic Meridional
 1280 Overturning Circulation to the melting from northern glaciers in climate change
 1281 experiments. *Geophys. Res. Lett.* 33, L07711. doi:10.1029/2006GL025765.
 1282 Sy, A., M. Rhein, J.R.N. Lazier, K.P. Koltermann, J. Meincke, A. Putzka, M. Bersch (1997),
 1283 Surprisingly rapid spreading of newly formed intermediate waters across the North
 1284 Atlantic Ocean. *Nature* 386, 675–679.
 1285 Talley, L.D. and M.S. McCartney (1982), Distribution and circulation of Labrador Sea
 1286 Water. *J. Phys. Oceanogr.*, 12(11), 1189-1205.

- Tang, C.C.L., C.K. Ross, T. Yao, B. Petrie, B.M. DeTracey, and E. Dunlap (2004), The circulation, water masses and sea-ice of Baffin Bay, *Progress in Oceanography*, 63, 183-228.
- Vage, K., R.S. Pickart, V. Thierry, G. Reverdin, C. M. Lee, B. Petrie, T.A. Agnew, A. Wong, and M.H. Ribergaard (2009), Surprising return of deep convection to the subpolar North Atlantic Ocean in winter 2007-2008, *Nature Geoscience*, 2(1), 67-72.
- Velicogna, I. (2009), Increasing rates of ice mass loss from the Greenland and Antarctic ice sheets revealed by GRACE. *Geophys. Res. Lett.* 36, L19503. doi:10.1029/2009GL040222.
- Vested, H.J., P. Justesen, and L. Ekabjaerg (1992), Advection-dispersion modeling in three dimensions, *Appl. Math. Modelling*, 16, 506-519.
- Vizcaino, M., et al., (2008), Longterm ice sheet-climate interactions under anthropogenic greenhouse forcing simulated with a complex Earth System Model. *Clim. Dyn.* 31, 665-690. doi:10.1007/s00382-008-0368-7.
- Watson, A.J., M.-J. Messias, E. Fogelqvist, K.A. Van Scoy, T. Johannessen, K.I.C. Oliver, D.P. Stevens, F. Rey, T. Tanhua, K.A. Olsson, F. Carse, K. Simonsen, J.R. Ledwell, E. Jansen, D.J. Cooper, J.A. Kruepke, and E. Guilyardi (1999), Mixing and convection in the Greenland Sea from a tracer-release experiment, *Nature*, 401, 902-904.
- Walczowski, W., *Atlantic Water in the Nordic Seas. Properties, Variability, Climatic Importance*. Springer International Publishing, 174 pp.
- Walczowski, W., and J. Piechura (2006), New evidence of warming propagating toward the Arctic Ocean, *Geophys. Res. Lett.*, 33, L12601, doi:10.1029/2006GL025872.
- Yashayaev, I. (2007), Hydrographic changes in the Labrador Sea, 1960-2005, *Progress in Oceanography* 73, 242-276.
- Yashayaev, I. and J. W. Loder (2009), Enhanced production of Labrador Sea Water in 2008, *Geophys. Res. Lett.*, 36, L01606, doi:10.1029/2008GL036162.
- Yashayaev, I. and D. Seidov (2015), The role of the Atlantic Water in multidecadal ocean variability in the Nordic and Barents Seas, *Progress in Oceanography*, 132, 68-127.
- Yashayaev, I., D. Seidov and E. Demirov (2015), A new collective view of oceanography of the Arctic and North Atlantic basins, *Progress in oceanography*, 132, 1-21
- Zweng, M.M. and A. Munchow (2006), Warming and freshening of Baffin Bay, 1916-2003, *JGR*, 111, C07016, doi: 10.1029/2005JC003093.

Figure Captions

Figure 1. Map of the sub-Arctic seas with general circulation and geographic names shown, including the regions selected for tracer budget analysis (section 4). Three main basins include the Nordic Seas (NS), the Labrador Sea (LS), and Baffin Bay (BB). The boxes are interior regions in the Greenland Sea (GS), Iceland Sea (IS), and interior Labrador Sea (IL). The numbers indicate segments bounding the regions. Color coding is used for ease of identification of the segments bounding the regions. The dashed black line across Davis Strait shows the location of the 2004-present moored array. Abbreviated currents: CSC – Continental Slope Current, FC – Faroe Current, SC – Shetland Current, NCC – Norwegian Coastal Current, NwAC – Norwegian Atlantic Current, WSC – West Spitsbergen Current, EGC – East Greenland Current, JMC – Jan Mayen Current, EIC – East Icelandic Current, NIC – North Icelandic Current, WGC – West Greenland Current, BIC – Baffin Island Current, LC – Labrador Current.

Figure 2. Seasonal and interannual variability of Greenland freshwater fluxes from *Bamber et al.*, [2012], 1990-2010. The fluxes include both solid and liquid discharge. (a) Locations and volume fluxes ($\text{km}^3 \text{ mo}^{-1}$) of Greenland freshwater sources along the coast. The bar diagrams show monthly averaged Greenland freshwater fluxes between 1990-2010 ($\text{km}^3 \text{ mo}^{-1}$) integrated over the regions delineated with the black lines. At the upper right, monthly runoff integrated over the whole Greenland coast is shown. The vertical black lines on top of the bars indicate the range of the monthly runoff during 1990-2010. Long-term (1990-2010) mean runoff ($\text{km}^3 \text{ yr}^{-1}$) is listed for every region. (b) Stacked bar diagram of the annual Greenland freshwater flux in the regions. The black line depicts the total annual freshwater flux.

Figure 3. Mean salinity and salinity anomalies derived from the observational program in Davis Strait data between 2004 and 2013 (the location of the moored array is shown in Figure 1). (a) Mean salinity. The major water masses are indicated: ArW – Arctic Water, TrW – Transitional Water, WGIW – West Greenland Irminger Water, WGSW – West Greenland Shelf Water. (b) Annual salinity anomalies (ΔS) relative to the 2004–2013 mean for the individual water masses. The other water mass refers to the near surface water that is too warm to be classified as ArW or is near surface water along the West Greenland slope right at the shelf that is too salty to be WGSW and too fresh to be WGIW. (c) Annual salinity anomalies relative to the 2003-2013 mean. (d) Time vs latitude diagram showing salinity anomalies along the section shown with the red line on the map to the left for the time period from 1990 to 2014 relative to the 1960-1990 monthly climatology. Salinity climatology and

anomalies are derived from the Met Office Hadley Centre subsurface ocean salinity dataset (EN.4.1.1).

Figure 4. Time-depth diagrams of salinity in the central Greenland Sea (a) and Iceland Sea (b) from hydrographic observations in boxes GS and IS shown in Figure 1. Hydrographic measurements (salinity, temperature, and pressure from bottle and CTD data) in the Greenland and Iceland seas from 1950 to 2010 are from the ICES Dataset on Ocean Hydrography (www.ocean.ices.dk) and the Pangaea database (www.pangaea.de). Note a very shallow (upper 25 m) freshening signal at both locations.

Figure 5. Daily mean tracer concentration in the first model layer ($\sim 3\text{m}$) on December 17th for every model year from AO-HYCOM. The concentration is on the natural-log scale (kg m^{-3}).

Figure 6. Five-day mean tracer concentration in the surface model layer ($\sim 1\text{m}$) for output including December 17th for every model year from NEMO-LIM2. The concentration is on the natural-log scale (kg m^{-3}).

Figure 7. Daily mean tracer concentration in the surface model layer ($\sim 2\text{m}$) on December 17th for every model year from ICMMG. The concentration is on the natural-log scale (kg m^{-3}).

Figure 8. Tracer budget for the Nordic Seas region. (a) Volume-integrated tracer content (kg). Colors designate different models. (b—d) 60-day low-pass filtered time series of the tracer fluxes (kg s^{-1}) across the sections from the model experiments. Colors designate fluxes across individual sections shown in the inset in (b). Section numbers correspond to Figure 1. Note the different scale in 8b vs 8c and 8d.

Figure 9. Same as Figure 8 but for the Greenland Sea region. Note the different scales in 9b–9d.

Figure 10. Same as Figure 8 but for the Iceland Sea region. Note the different scales in 10b–10d.

Figure 11. Same as Figure 8 but for the Labrador Sea region.

Figure 12. Same as Figure 8 but for the interior Labrador Sea region.

Figure 13. Same as Figure 8 but for Baffin Bay. (d) In ICMMG, section 22 is over land, thus the flux is not shown.

Figure 14. Mass fraction (%) of the volume-integrated tracer in 3 regions (NS, LS, and BB) relative to the total tracer mass integrated over the model domain (limited by $\sim 38^\circ\text{N}$ for NEMO-LIM2 and ICMMG). (a) AO-HYCOM; (b) NEMO-LIM2; (c) ICMMG. Shown are mass fraction values after 3, 6, 9, 12, and 14 years of the simulation. The colored bars inside

the “Nordic Seas” and “Labrador Sea” illustrate the mass fraction of the tracer integrated over the interior boxes (blue - GS, red - IS, and green - IL) relative to the total tracer mass (the values are listed in colored numbers). Also listed is the total mass fraction of the tracer within the three regions.

Figure 15. Distribution of the tracer concentration (kg m^{-3}) along the vertical section (g) from the numerical experiments on December 17th after 7 and 14 years. In a–f, the vertical axis is depth in meters. (a, b) AO-HYCOM: black contours are interfaces of the vertical layers from a single model output time. (c) NEMO-LIM2. (e, f) ICMMG. (g) Section lines. The red numbers on the blue line correspond to the distances (km) along the line also shown on the horizontal axis on the diagrams.

Figure 16. Mean (top row) and eddy (bottom row) tracer fluxes ($\text{kg s}^{-1} \text{ m}^{-1}$) integrated over the upper 150 m calculated from daily-mean fields (a, b) AO-HYCOM, (c, d) NEMO-LIM2, and (e, f) ICMMG for year 9. The fluxes are one-year-averaged.

Figure 17. Mass gain (time integrated net tracer flux, kg) from the mean tracer flux (blue) and eddy tracer flux (red) into the interior convective regions (Figure 1) during years 8–12 of the simulations from (a–c) AO-HYCOM, (d–f) NEMO-LIM2, and (g–i) ICMMG. The horizontal axis is time (model years).

Figure 18. Hydrographic changes in the sub-Arctic regions (in rows) from AO-HYCOM. (a) Time series of freshwater content (black) and salinity (blue) in the upper 400 m during the simulation. (b) Salinity change of the upper 400m vs cumulative time-integrated Greenland freshwater flux. Linear regression parameters (Eq. 5) and coefficient of determination are listed. (c) Estimated salinity change in the upper 400m for time-integrated anomaly of Greenland freshwater flux for 1990-2010.

1413 **Figures**
1414
1415